

<p>(51) International Patent Classification 4 : C12N 15/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 89/ 01038</p> <p>(43) International Publication Date: 9 February 1989 (09.02.89)</p>
<p>(21) International Application Number: PCT/US88/02445</p> <p>(22) International Filing Date: 20 July 1988 (20.07.88)</p> <p>(31) Priority Application Number: 077,188</p> <p>(32) Priority Date: 24 July 1987 (24.07.87)</p> <p>(33) Priority Country: US</p> <p>(71) Applicant: CETUS CORPORATION [US/US]; 1400 Fifty-Third Street, Emeryville, CA 94608 (US).</p> <p>(72) Inventors: KAWASAKI, Ernest, Seigo ; 2601 San Mateo Street, Richmond, CA 94804 (US). DEVLIN, James, Joseph ; 522 Silverado Drive, Lafayette, CA 94549 (US). MARTIN, George ; 2227 McKinley Avenue, Berkeley, CA 94703 (US). O'ROURKE, Edward ; 111 Temescal Circle, Emeryville, CA 94608 (US). CLARK, Robin ; 3736 Woodruff Avenue, Oakland, CA 94602 (US).</p>	<p>(74) Agent: HALLUIN, Albert, P.; Cetus Corporation, 1400 Fifty-Third Street, Emeryville, CA 94608 (US).</p> <p>(81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), DK, FI, FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), NO, SE (European patent).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: PRODUCTION OF BIOLOGICALLY ACTIVE FORMS OF CSF USING A BACULOVIRUS (AcNPV)-INSECT CELL EXPRESSION SYSTEM</p>		
<p>(57) Abstract</p> <p>The expression of biologically active CSFs in insect cells (<i>Spodoptera frugiperda</i>) is disclosed. Numerous forms of CSF-I are expressed and secreted as biologically active dimers. DNA transfer vectors and expression vectors that encode forms of CSF-I and allow for their expression are disclosed and claimed. G-CSF is additionally expressed in insect cells in active form. DNA transfer vectors and expression vectors that encode G-CSF are disclosed and claimed.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

PRODUCTION OF BIOLOGICALLY ACTIVE
FORMS OF CSF USING A BACULOVIRUS (AcNPV)-
INSECT CELL EXPRESSION SYSTEM

The present invention relates to the field of molecular
5 biology and proteins. In particular, the invention relates to the
achievement of high levels of expression of biologically active colony
stimulating factors using an insect cell/baculovirus expression
system.

10 Numerous different expression systems are now available to
the genetic engineer who wishes to express a cloned gene of
interest. Generally the goal is the production of high levels of
biologically active material. The requirements for the expression
system chosen to accomplish this goal are dependent upon the nature of
the protein to be expressed.

15 The value of utilizing prokaryotic host vector systems for
the synthesis of desirable eukaryotic proteins is diminished by
certain limitations inherent in such systems. For instance, the mRNA
transcript or protein product of such systems may be unstable in the
prokaryote. In addition, before a protein will be synthesized within
20 a prokaryotic cell, the DNA sequence introduced into the microorganism
must be free of intervening DNA sequences, nonsense sequences, and
initial or terminal sequences which encode for polypeptide sequences
which do not comprise the active eukaryotic protein. Further, some
eukaryotic proteins require modification after synthesis (e.g.,
25 glycosylation and all membrane associated processing) to become
biologically active, and prokaryotic cells are generally incapable of
such modifications.

30 Various nonviral eukaryotic host vector systems are also
available for the expression of heterologous proteins. Certain
limitations are inherent in each of these systems as well. For
example, high levels of expression are frequently difficult to obtain
in yeast systems where autonomously replicating vectors may be
unstable. Additionally, glycosylation patterns in yeast differ from
those in higher eukaryotes.

Limitations encountered with mammalian host vector systems include difficulties in host cell culturing and its scale-up. The expense of mammalian cell culture media as well as a requirement for serum often restricts its use on a large scale and complicates the use of these systems for production of parenterally administered pharmaceuticals. Furthermore, levels of expression in these systems are generally substantially lower than that obtained in prokaryotic or viral expression systems.

The use of viruses in eukaryotic host-vector systems has been the subject of a considerable amount of recent investigation and speculation. However, some viral vector systems also suffer from significant disadvantages and limitations which diminish their utility. For example, a number of eukaryotic viral vectors are either tumorigenic or oncogenic in mammalian systems and create potentially health and safety problems associated with resultant gene products and accidental infection.

The baculovirus expression vector system involved in the instant invention overcomes many of the above-mentioned limitations. Baculoviruses are insect pathogenic viruses which, until recently, were studied mostly for their potential use as viral insecticides for control of agriculturally important insect pests. Because certain baculoviruses are highly virulent for pest insects, some of the most promising have been commercially developed and are used as biological pesticides (Miltnerburger and Krieg 1984 Bioinsecticides:II: Baculoviridae. Adv. Biotechnol. Processes 3:291; Granados, R.R. and Federici, B.A. eds. The Biol. of Baculoviruses Vol II, Boca Raton, FL: CRC Press, Inc. 1986). Baculoviruses are very stable and are able to persist for longer times in the environment than other animal viruses. This unusual biological stability is the result of a unique association of the infectious virus particles and a viral occlusion which is a crystalline assembly of a viral encoded structural protein called polyhedrin. Late in viral replication, baculovirus particles become embedded in a protein occlusion composed of the polyhedrin protein. Insects acquire a baculovirus disease by ingesting these occluded virus (OV) which contaminate their food supply. The

polyhedrin matrix protects the virus particles in the environment and during their passage through the foregut of the insect. In the insect midgut, the alkaline pH activates the dissolution of the polyhedrin crystalline matrix resulting in the release of many viruses. The virus become absorbed by the midgut epithelial cells and initiate the infection process.

There is a second infectious form of nuclear polyhedrosis viruses (NPVs), known as the extracellular or nonoccluded virus (NOV) form, which is generated by the budding of viral nucleocapsids through the plasma membrane of the infected cells. NOVs are responsible for spreading a secondary infection via the hemolymph of the insect. It is the NOV form of the virus which is infectious in insect cell cultures; the occluded (OV) form is not infectious in cell cultures since dissolution of the crystalline matrix occurs only at high alkaline pH (i.e., pH 10.5).

The formation of NOVs and OVs occurs in a biphasic manner during the infection process. NOVs are abundantly produced before occlusion is initiated. During a typical synchronous infection of fully permissive cell lines, the majority of NOVs are produced between 12 and 24 hr post-infection (p.i.). The synthesis of polyhedrin is initiated at 20 hr p.i. and does not reach maximal levels until 48 to 72 hr p.i. The significance of this temporal regulation with respect to the expression vector system is that foreign gene products that may have adverse effects on the cell should not diminish the production of progeny NOVs to be used for further infection.

Of the 450-500 species of known baculoviruses, practically all encode for a polyhedrin protein. As previously discussed, the viral occlusion is a paracrystalline assembly of a polyhedrin monomer which, for most viruses, has an average molecular weight of 28,000-30,000 daltons (Summers, M.D. and Smith, G.E., 1978 Virology 84:390). Baculoviruses are unique among animal viruses, not only in the protective function of the viral occlusion in the viral life cycle but also because the polyhedrin gene is the most highly expressed eucaryotic virus gene known. The polyhedrin protein can accumulate to

greater than 1 mg/ml of infected cultured insect cells (70-75% of the total cellular protein) or can comprise up to 25% of the total protein of an infected insect. Although very highly expressed, neither the polyhedrin gene nor its protein is essential for viral infection or replication in cultured insect cells or insects, thus making the polyhedrin gene an ideal target for genetic manipulation.

The most extensively studied baculovirus is the Autographa californica nuclear polyhedrosis virus (AcNPV). The Autographa californica host for AcNPV is a moth commonly referred to as the alfalfa looper. Studies of the physical and functional organization of the AcNPV genome have resulted in the mapping, cloning, and sequencing of the AcNPV polyhedrin gene and its regulatory sequences (Iddekinge et al. (1983) Virology 131:561; Smith et al. (1983) J. Virol. 46:584). Not only does the polyhedrin gene exhibit a strong promoter, but expression can continue late in infection well beyond the point of repression of nearly all other baculovirus and host genes.

The genetic engineering of the baculovirus polyhedrin gene for high level expression of a heterologous protein, in this case recombinant human beta-interferon was first reported by Smith et al. Mol. Cell. Biol. 3(12):2156-2165 (1983)). Since then, human interleukin-2 has been expressed in insect cells by a baculovirus expression vector as described by Smith et al. (Proc. Natl. Acad. Sci. USA 32:8404-8408 (1985)). Recently, the synthesis of functional human T-cell leukemia virus Type I p40^X protein using a baculovirus expression vector has been reported (Jeang, K.T., et al., J. Virol. 61:708-713 (1987)). Other heterologous proteins that have been expressed in this system are summarized in Summers et al. ("Genetic Engineering of the Genome of the Autographa californica Nuclear Polyhedrosis Virus" Banbury Report: Genetically Altered Viruses in the Environment, 22:319-329, Cold Spring Harbor Laboratory (1985)).

The baculovirus expression system has several advantages for the expression of foreign genes in comparison to other prokaryotic, yeast or mammalian cell expression vector systems. First, high levels

of expressed proteins are possible. Greater than 1.0 mg per ml of polyhedrin protein is normally produced in infected cells. Another non-essential occlusion-related viral protein, p10, is also abundantly produced and its promoter has reportedly been used to drive foreign gene expression (D.W. Miller et al. Genetic Engineering Principles & Methods 8:277-298, Setlow and Hollaender, eds. New York: Plenum Press, 1986). Using the polyhedrin gene promoter, heterologous gene expression levels never reach polyhedrin levels but are usually in the range of tens to hundreds of micrograms per ml (Summers et al. (1985), p. 321, supra). Secondly, in contrast to those produced in bacterial or yeast cells, recombinant proteins produced in insect cells may be co- and post-translationally processed in a manner similar to what occurs in mammalian cells. In at least one case, glycosylation of IFN-beta in infected insect cells has been reported (G.E. Smith et al. (1983) supra). Whereas about 40% of the natural IL-2 produced in human Jurkat cells is not glycosylated, there was no evidence of any glycosylation of the recombinant IL-2 produced in insect cells (G.E. Smith, et al., (1985), supra). In addition, correct cleavage of mammalian secretory signal peptides has been observed (G.E. Smith et al., (1983), supra; G.E. Smith et al., (1985), supra; D.W. Miller et al., (1986), supra).

As described by M. B. Ladner et al., (EMBO J. (1987) 6:2683-2698 and incorporated herein by reference, the production of biologically active CSF-1 is complicated by its high degree of posttranslational processing which includes glycosylation and dimerization. In addition there are a large number of cysteine residues, in particular, in the N-terminal portion of the protein. There are, in fact, a total of ten cysteine residues in the "long form" of CSF and seven in the "short form". Both thus contain cysteine residues at positions 7, 31, 48, 90, 102, 139, 146, the long form has additional cysteines at 157, 159 and 225. It is believed that processing to form dimer includes formation of multiple intrachain and at least one interchain disulfide bond(s). While prokaryotic systems may be employed for CSF-1, refolding protocols are necessary to produce high yields of biologically active product. It

would be desirable to have a high level expression system that produces biologically active dimers of CSF-1.

As described in PCT Publication No. W088/01297 published February 25, 1988, and incorporated herein by reference, the native DNA sequence encoding G-CSF has not proved to be effective for the production of high levels of G-CSF in prokaryotic hosts. Not only does the insect cell expression system provide an intriguing alternative but the possibilities of achieving the appropriate post-translational processing and recovery of G-CSF from the culture medium would be advantageous in the production of pharmaceutically useful CSFs.

The invention relates, in one respect, to methods for producing by recombinant DNA technology biologically active CSFs and in the case of CSF-1, a dimerized form of CSF-1, using host insect cells infected with a recombinant baculovirus expression vector. Accordingly, one aspect of the invention relates to growing the infected insect cells under suitable conditions to produce the desired recombinant CSFs and recovering the biologically active polypeptide or dimerized polypeptide from the culture medium.

In another aspect, the invention is directed to recombinant baculovirus expression vectors which are capable of effecting the expression of CSFs, to the host cells infected with such vectors, and to cultures thereof.

One aspect of the invention concerns recombinant baculovirus expression vectors in which CSFs are expressed under the transcriptional control of a baculovirus promoter. In one aspect of the invention the baculovirus promoter is the polyhedrin gene promoter.

Yet another aspect of the invention relates to recombinant baculovirus expression vectors that are capable of expressing CSFs and which may encode either the long or short forms of CSF-1, muteins thereof, CSF-1 in which the amino terminal 4 codons are altered, or G-CSF.

Also, aspects of the invention are the recombinant baculovirus transfer vectors which are used to transfer the desired recombinant gene into the baculovirus genome.

Figure 1 shows the DNA sequence of the CSF-1 gene from pcCSF-17.

Figure 2 shows the DNA sequence of original cDNA clone number 4. The sequence is missing a few nucleotides at the 5' end compared to the LCSF sequence in pcDBhuCSF-4 which was completed by substitution of upstream codons from pcCSF-17 (see Example 2, Section A.2.)

Figure 3 shows the DNA sequence of G-CSF from pP12.

Figure 4 shows the DNA sequence comparison between the recombinant baculovirus transfer vectors pAcC1-C5. The carrots indicate restriction endonuclease cleavage sites.

Figure 5 shows the DNA sequence comparison for the regions around the CSF-1 translational start codon between the recombinant baculovirus transfer vectors pAcM1, pAcM2, pAcM3, pAcM4, pAcM6, pAcM10 and the polyhedrin gene.

Figure 6 shows SDS-PAGE analysis of immunoprecipitated baculovirus expression system products.

Figure 7 shows Western blot analysis of G-CSF expressed by the baculovirus system.

Figure 8 shows expression levels of various recombinant polypeptides in the baculovirus expression vector system.

Modes for Carrying Out the Invention

A. Definitions

"Colony stimulating factor-1 (CSF-1)" refers to a protein which exhibits the spectrum of activity understood in the art for CSF-1, i.e., when applied to the standard in vitro colony stimulating assay of Metcalf, D., J. Cell Physiol. (1970) 76:89, it results in the

formation of primarily macrophage colonies. Native CSF-1 is a glycosylated dimer; dimerization may be necessary for activity. Contemplated within the scope of the invention and within the definition of CSF-1 are both the dimeric and monomeric forms. The monomeric form may be converted to the dimeric form by in vitro provision of host intracellular conditions suitable for dimerization of the monomer.

There appears to be some species specificity: Human CSF-1 is operative both on human and on murine bone marrow cells; murine CSF-1 does not show activity with human cells. Therefore, "human" CSF-1 should be positive in the specific murine radioreceptor assay of Das, S. K., et al., Blood (1981) 58:630, although there is not necessarily a complete correlation. The biological activity of the protein will generally also be inhibited by neutralizing antiserum to human urinary CSF-1 (Das, S. K., et al., supra). However, in certain special circumstances (such as, for example, where a particular antibody preparation recognizes a CSF-1 epitope not essential for biological function, and which epitope is not present in the particular CSF-1 mutein being tested) this criterion may not be met.

Certain other properties of CSF-1 have been recognized more recently, including the ability of this protein to stimulate the secretion of series E prostaglandins, interleukin-1, and interferon from mature macrophages (Moore, R., et al., Science (1984) 223:178; Ralph, P. et al. (1986) Immunobiol., 172:194; Ralph, P. et al. (1987) Cell. Immunol., 105:270). The mechanism for these latter activities is not at present understood, and for purposes of definition herein, the criterion for fulfillment of the definition resides in the ability to stimulate the formation of monocyte/macrophage colonies using bone marrow cells from the appropriate species as starting materials, under most circumstances (see above) the inhibition of this activity by neutralizing antiserum against purified human urinary CSF-1, and, where appropriate for species type, a positive response to the radioreceptor assay. (It is known that the proliferative effect of CSF-1 is restricted to cells of mononuclear phagocytic lineage (Stanley, E. R., The Lymphokines (1981), Stewart, W. E., II, et al.,

ed., Humana Press, Clifton, NJ), pp. 102-132) and that receptors for CSF-1 are restricted to these cell types (Byrne, P. V., et al., Cell Biol. (1981) 91:848)).

Figure 1 shows the amino acid sequence for a particular form of human CSF-1 encoded by the recombinant cDNA clone pcCSF-1 (Kawasaki, E.S., et al. (1985), Science 230:291). This protein contains 224 amino acids in the mature sequence and a leader sequence of 32 amino acids. The protein produced as the expression product of this clone is active in assays specific for CSF-1, namely, the bone marrow proliferation assay (wherein the activity is destroyed by addition of anti-CSF-1 antibodies), colony stimulation assays, and a radioreceptor assay. Further characterization of the biological and molecular properties of human and murine csF-1 is disclosed by P. Ralph et al. in Molecular Basis of Lymphokine Action (D. R. Webb et al., eds., The HUMANA Press, Inc., 1987, pp. 295-311).

For convenience, the mature protein amino acid sequence of the monomer portion of a dimeric protein shown in Figure 1, deduced from the cDNA clone illustrated herein, is designated mCSF-1 (mature CSF-1). Figure 1 shows the presence of a 32 residue putative signal sequence, which is presumably cleaved upon secretion from mammalian cells; mCSF-1 is represented by amino acids 1-224 shown in that figure. Specifically included in the definition of human CSF-1 are mutants which monomers and dimers are mCSF-1 and related forms of mCSF-1. CSF-1 derived from other species may fit the definition of "human" CSF-1 by virtue of its display of the requisite pattern of activity as set forth above with regard to human substrate.

Also for convenience, the amino acid sequence of mCSF-1 will be used as a reference and other sequences which are substantially equivalent to this in terms of CSF-1 activity will be designated by referring to the sequence shown in Figure 1. The substitution of a particular amino acid will be noted by reference to the amino acid residue which it replaces. Thus, for example, ser₉₀CSF-1 refers to the protein which has the sequence shown in Figure 1 except that the amino acid at position 90 is serine rather than cysteine. Numerous

5 muteins have been described in commonly owned PCT Publication Nos. W086/04607 published August 14, 1986 and PCT Publication No. W088/03173 published May 5, 1988 each incorporated herein by reference and are included within the definition of CSF-1 according to the invention so long as the biological activity of CSF-1 is maintained. It is expected, of course, that such modifications may quantitatively or qualitatively affect the activity, either by enhancing or diminishing the activity of the protein in various assays.

10 Additional CSF-1 proteins, which have the same colony stimulating activity described in Metcalf supra, contain substantially longer amino acid sequences deduced from the recovered cDNA, which include a 298 amino acid "extra" segment at approximately residue 150 of the pcCSF-17 encoded protein and expression of this cDNA results in CSF-1 activity in these specific CSF-1 assays, as well.

15 For convenience, the CSF-1 encoded by pcCSF-17 will be referred to as the "short form" of the protein, and for consistency with previous designations, as "mature" or mCSF-1. The 522 amino acid sequence encoded in, for example, pcDBhuCSF-4 and its corresponding clones will be referred to as the "long form", or LCSF-1. This sequence is shown in Figure 2. LCSF-1 is further described in PCT Publication No. W087/06954 published November 19, 1987, and in commonly owned PCT Publication No. W088/03173 published May 5, 1988. The abbreviations used herein also include "huCSF-1" for all human forms of the protein.

25 "G-CSF" as used herein means a protein having the effect of stimulating the production of primarily granulocyte colonies or granulocyte-macrophage colonies in a colony forming assay using bone marrow cell progenitors of an appropriate species. A protein having this activity has the deduced amino acid sequence shown in Figure 3 as disclosed in commonly owned EP Publication No. 256,843 published February 24, 1988, and in PCT Publication No. W088/01297 published February 25, 1988 incorporated herein by reference.

30 G-CSF may be isolated from the MIA PaCa-2 cell line. In addition, G-CSF specific RNA sequences are clearly detectable in the mRNA of LD-1 cells and 5637 cells. Thus, G-CSF obtained by cloning

mRNA of LD-1 cells and 5637 cells. Thus, G-CSF obtained by cloning from these cell lines as well as any other cell line which is G-CSF probe positive, is considered within the scope of the invention. Mature G-CSF produced by the LD-1 cell line has been isolated and amino-acid sequenced. 65% of the LD-1 produced protein has a Thr residue at the NH₂-terminus and 35% has a Pro residue at the NH₂-terminus. Further disclosure on the cloning and expression of G-CSF are found in commonly owned PCT Publication No. W088/01297 published February 25, 1988 and EP Publication No. 256,843 (supra).

The mature G-CSF protein amino acid sequence shown in Figure 3, deduced from the cDNA clone, is designated mG-CSF (mature G-CSF) beginning at the amino acid residue threonine designated +1. There is a 30 residue putative signal sequence, which is presumably cleaved upon secretion from mammalian cells; mG-CSF is represented by amino acids 1-174 shown in that figure. Specifically included in the definition of human G-CSF are muteins which monomers and dimers, if any, are G-CSF and related forms of G-CSF, designated by their differences from mG-CSF. G-CSF derived from other species may fit the definition of "human" G-CSF by virtue of its display of the requisite pattern of activity as set forth above with regard to human substrate.

Also for convenience, the amino acid sequence of G-CSF will be used as a reference and other sequences which are substantially equivalent to this in terms of G-CSF activity will be designated by referring to the sequence shown in Figure 3. The substitution of a particular amino acid will be noted by reference to the number of the amino acid residue which it replaced. Thus, for example, ser₆₀ G-CSF refers to the protein which has the sequence shown except that the amino acid at position 60 is serine rather than proline.

As is the case for all proteins, the precise chemical structure of CSF-1 or G-CSF depends on a number of factors. As ionizable amino and carboxyl groups are present in the molecule, a particular protein may be obtained as an acidic or basic salt, or in neutral form. All such preparations which retain their activity when placed in suitable environmental conditions are included in the definition. Further, the primary amino acid sequence may be augmented

by derivatization using sugar moieties (glycosylation) or by other supplementary molecules such as lipids, phosphate, acetyl groups and the like, more commonly by conjugation with saccharides. The primary amino acid structure may also aggregate to form complexes. Certain aspects of such augmentation are accomplished through post-translational processing systems of the producing host; other such modification may be introduced in vitro. In any event, such modifications are included in the definition so long as the activity of the protein, as defined above, is not destroyed. It is expected, of course, that such modifications may quantitatively or qualitatively affect the activity, either by enhancing or diminishing the activity of the protein in the various assays. Further, individual amino acid residues in the chain may be modified by oxidation, reduction, or other derivatization, and the protein may be cleaved to obtain fragments which retain activity. Such alterations which do not destroy activity do not remove the protein sequence from the definition.

Modifications to the primary structure itself by deletion, addition, or alteration of the amino acids incorporated into the sequence during translation can be made without destroying the activity of the protein. Such substitutions or other alterations result in proteins having an amino acid sequence which falls within the definition of proteins "having an amino acid sequence substantially equivalent to that of CSF-1 or G-CSF."

"Operably linked" refers to juxtaposition such that the normal function of the components can be performed. Thus, a coding sequence "operably linked" to control sequences refers to a configuration wherein the coding sequence can be expressed under the control of these sequences.

"Control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. Eukaryotic cells including the insect cells of the instant invention are known to utilize promoters and polyadenylation signals.

"Expression system" refers to DNA sequences containing a desired coding sequence and control sequences in operable linkage, so that hosts transformed with these sequences are capable of producing the encoded proteins. These DNA sequences may also direct the synthesis of the encoded proteins in an in vitro cellular environment. In order to effect transformation, the expression system may be included on a transfer vector; however, the relevant DNA may then also be integrated into the viral chromosome to result in a recombinant viral genome.

As used herein "cell", "cell line", and "cell culture" are used interchangeably and all such designations include progeny. Thus "transformants" or "transformed cells" includes the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Mutant progeny which have the same functionality as screened for in the originally transformed cell are included. Where distinct designations are intended, it will be clear from the context.

"Infection" as used herein refers to the invasion of cells by pathogenic viral agents where conditions are favorable for their replication and growth. "Transfection" refers to a technique for infecting cells with purified nucleic acids by adding calcium chloride to solutions of DNA containing phosphate or other appropriate agents such as dextran sulfate thereby causing the DNA to precipitate and be taken up into the cells.

"Recombinant transfer vector" refers to a plasmid containing a "heterologous" gene under the control of a functional promoter (e.g., polyhedrin or p10 promoter) and flanked by viral sequences. The "recombinant expression vector" is formed after cotransfection of the recombinant transfer vector and wild-type baculovirus DNA into host insect cells whereupon homologous recombination occurs between the viral sequences flanking the heterologous gene and the homologous sequences in the wild-type viral DNA. This results in the replacement of wild-type sequences in the virus with the transfer vector sequences

between the crossover points. The recombinant expression vector is the recombinant viral DNA containing the desired heterologous gene.

B. General Description

Interest in the growth and differentiation of granulocytes and macrophages from bone marrow progenitors has prompted the study of the colony-stimulating factors (CSFs) that regulate these processes. The subject has been recently reviewed by D. Metcalf in Blood, 67:257 (1986). One of these factors, termed macrophage-CSF (M-CSF or CSF-1) stimulates the in vitro growth of predominantly macrophage colonies from bone marrow stem cells. Another of these factors, termed granulocyte-CSF (G-CSF) stimulates the in vitro growth of predominantly granulocyte colonies from bone marrow stem cells.

The CSF-1 proteins of the invention are capable both of stimulating monocyte-precursor-macrophage cell production from progenitor marrow cells, thus enhancing the effectiveness of the immune system, and of stimulating such functions of these differentiated cells as the secretion of lymphokines in the mature macrophages.

In one application, these proteins are useful as adjuncts to chemotherapy. It is well understood that chemotherapeutic treatment results in suppression of the immune system. Often, although successful in destroying the tumor cells against which they are directed, chemotherapeutic treatments result in the death of the subject due to this side effect of the chemotoxic agents on the cells of the immune system. Administration of CSF-1 to such patients, because of the ability of CSF-1 to mediate and enhance the growth and differentiation of bone marrow-derived precursors into macrophages, results in a restimulation of the immune system to prevent this side effect, and thus to prevent the propensity of the patient to succumb to secondary infection. Other patients who would be helped by such treatment include those being treated for leukemia through bone marrow transplants; they are often in an immunosuppressed state to prevent rejection. For these patients also, the immunosuppression could be reversed by administration of CSF-1.

In addition to the cDNA encoding the 224 amino acid CSF-1 monomer described above, cDNAs from both human and murine systems have been obtained which include DNA encoding an additional 298 amino acid sequence. These cDNAs express to produce "long form" CSF-1 proteins which fulfill the criteria for biological activity of this protein.

In general outline, the human cDNA encoding the 224 amino acid form of the protein was used as a probe to recover the sequences encoding murine CSF-1 from a cDNA bank prepared in lambda gt10 from L-929 mRNA which had been enriched for CSF-1 production capability. Two clones were recovered which encode similar 520 amino acid proteins. The clones diverge dramatically in the 3' untranslated region. One of the 3' untranslated regions is more than 2 kb and bears little resemblance to the corresponding human sequences; the other, shorter clone contains approximately 500 bp in the untranslated region and shows considerable homology to the corresponding human DNA.

The longer forms of CSF-1 obtained from the murine library were then used as a basis to prepare probes to retrieve longer human sequences, if any. Based on comparison of the murine cDNAs to the human genomic sequence, a region of the gene previously thought to lie in an intron region putatively encodes an amino acid sequence showing considerable homology to the "extra" 298 amino acid segment contained in the murine sequence. This permitted construction of an oligonucleotide probe based on the "extra" DNA which had been, in the murine system, translated to protein. However, since the human genomic sequence was available, the probe was designed to accommodate the precise human sequence.

While pcCSF-17 had been prepared from MIA PaCa mRNA enriched for CSF-1-encoding materials, a subsequent cDNA library was prepared from total mRNA extracted from MIA PaCa cells and cloned into gt10. The gt10 library was first screened using pcCSF-17 sequences as probe, and selected probe-positive candidates were screened using an oligonucleotide probe based on the "extra" translated sequence of the murine cDNA, but modified to correspond to the related region in the human genome. Several clones encoding a corresponding "long form" of

a human protein were also obtained. Of course, the availability of DNA encoding each of these sequences provides the opportunity to modify the codon sequence so as to generate mutein forms also having CSF-1 activity.

5 For example, glycosylation sites within the coding sequence may be predicted by the presence of the sequence Asn. x. Ser or Thr which is found at glycosylated Asn residues in all known glycoproteins (Hughes, R.C. (1983) Glycoproteins New York: Chapman and Hall, p. 42).. The Asn residue within this sequence, for example, may be
10 substituted by site specific mutagenesis to remove the glycosylation site. As disclosed herein, two such substitutions were carried out in the CSF-1 molecule.

C. Methods Employed

C.1. Transformations and Transfections

15 Transformation of E. coli cells was done according to procedures set forth in T. Maniatis, E.F. Fritsch and J. Sambrook Molecular Cloning: A Laboratory Manual (1982) Cold Spring Harbor Press.

20 Transfections of Sf9 cells are accomplished using a modification of the calcium phosphate precipitation technique (Graham, F.L. et al., 1973, Virology 52:456) as modified for insect cells (Burand, J.P. et al. (1980), Virol., 101:286; Carstens, E.B. et al. (1980) Virol., 101:311) and further described by Summers, M.D. and Smith, G.E. (A Manual of Methods for Baculovirus Vectors and Insect
25 Cell Culture Procedures, Texas A & M Press: 1986)

C.2. Probing mRNA by Northern Blot; Probe of cDNA or Genomic Libraries

30 RNA is fractionated for Northern blot by agarose slab gel electrophoresis under fully denaturing conditions using formaldehyde (Maniatis, T., et al., supra pp 202-203) or 10 mM methyl mercury (CH₃HgOH) (Bailey, J. M., et al., Anal. Biochem. (1976) 70:75-85; and Sehgal, P. B., et al., Nature (1980) 288:95-97) as the denaturant.

For methyl mercury gels, 1.5% gels are prepared by melting agarose in running buffer (100 mM boric acid, 6 mM sodium borate, 10 mM sodium sulfate, 1 mM EDTA, pH 8.2), cooling to 60°C and adding 1/100 volume of 1 M CH₃HgOH. The RNA is dissolved in 0.5 x running buffer and denatured by incubation in 10 mM methyl mercury for 10 minutes at room temperature. Glycerol (20%) and bromophenol blue (0.05%) are added for loading the samples. Samples are electrophoresed for 500-600 volt-hr with recirculation of the buffer. After electrophoresis, the gel is washed for 40 minutes in 10 mM 2-mercaptoethanol to detoxify the methyl mercury, and Northern blots prepared by transferring the RNA from the gel to a membrane filter.

cDNA or genomic libraries are screened using the colony or plaque hybridization procedure. Bacterial colonies, or the plaques for phage are lifted onto duplicate nitrocellulose filter papers (S & S type BA-85). The plaques or colonies are lysed and DNA is fixed to the filter by sequential treatment for five minutes with 500 mM NaOH, 1.5 M NaCl. The filters are washed twice for five minutes each time with 5 x standard saline citrate (SSC) and are air dried and baked at 80°C for two hours.

The gels for Northern blot or the duplicate filters for cDNA or genomic screening are prehybridized at 25-42°C for 6-8 hours with 10 ml per filter of DNA hybridization buffer without probe (0-50% formamide, 5-6 x SSC, pH 7.0, 5 x Denhardt's solution (polyvinylpyrrolidone, plus Ficoll and bovine serum albumin; 1 x = 0.02% of each), 20-50 mM sodium phosphate buffer at pH 7.0, 0.2% SDS, 20 µg/ml poly U (when probing cDNA), and 50 µg/ml denatured salmon sperm DNA). The samples are then hybridized by incubation at the appropriate temperature for about 24-36 hours using the hybridization buffer containing kinased probe (for oligomers). Longer cDNA or genomic fragment probes were labeled by nick translation or by primer extension.

The conditions of both prehybridization and hybridization depend on the stringency desired, and vary, for example, with probe length. For example, conditions for relatively long (e.g., more than

30-50 nucleotide) probes may employ a temperature of 42°C-55°C and hybridization buffer containing about 20%-50% formamide. For the lower stringencies needed for oligomeric probes of about 15 nucleotides, lower temperatures of about 25°C-42°C, and lower formamide concentrations (0%-20%) are employed. For longer probes, the filters may be washed, for example, four times for 30 minutes, each time at 40°C-55°C with 2 x SSC, 0.2% SDS and 50 mM sodium phosphate buffer at pH 7, then washed twice with 0.2 x SSC and 0.2% SDS, air dried, and are autoradiographed at -70°C for 2 to 3 days. Washing conditions are somewhat less harsh for shorter probes. Other conditions are known in the art that may provide the results disclosed herein.

C.3. Vector Construction

Construction of suitable vectors containing the desired coding and control sequences employs standard ligation and restriction techniques which are well understood in the art and are described in Maniatis, T. et al., supra. Isolated plasmids, DNA sequences, or synthesized oligonucleotides are cleaved, tailored, and religated in the form desired.

Site specific DNA cleavage is performed by treating with the suitable restriction enzyme (or enzymes) under conditions which are generally understood in the art, and the particulars of which are specified by the manufacturer of these commercially available restriction enzymes. See, e.g., New England Biolabs, Product Catalog. In general, about 1 µg of plasmid or DNA sequence is cleaved by one unit of enzyme in about 20 µl of buffer solution; in the examples herein, typically, an excess of restriction enzyme is used to insure complete digestion of the DNA substrate. Incubation times of about one hour to two hours at about 37°C are workable, although variations can be tolerated. After each incubation, protein is removed by extraction with phenol/chloroform, and may be followed by ether extraction, and the nucleic acid recovered from aqueous fractions by precipitation with ethanol. If desired, size separation of the cleaved fragments may be performed by polyacrylamide gel or

agarose gel electrophoresis using standard techniques. A general description of size separations is found in Methods in Enzymology (1980) 65:499-560.

Restriction cleaved fragments may be blunt ended by treating with the large fragment of E. coli DNA polymerase I (Klenow) in the presence of the four deoxynucleotide triphosphates (dNTPs) using incubation times of about 15 to 25 minutes at 20 to 25°C in 50 mM Tris pH 7.6, 50 mM NaCl, 6 mM MgCl₂, 6 mM dTT and 5-10 μ M dNTPs. The Klenow fragment fills in at 5' sticky ends but chews back protruding 3' single strands, even though the four dNTPs are present. If desired, selective repair can be performed by supplying only one of the, or selected, dNTPs within the limitations dictated by the nature of the sticky ends. After treatment with Klenow, the mixture is extracted with phenol/chloroform and ethanol precipitated. Treatment under appropriate conditions with S1 nuclease results in hydrolysis of any single-stranded portion.

Synthetic oligonucleotides may be prepared by the triester method of Matteucci, et al., J. Am. Chem. Soc. (1981) 103:3185-3191 or using automated synthesis methods. Kinasing of single strands prior to annealing or for labeling is achieved using an excess, e.g., approximately 10 units of polynucleotide kinase to 1 nM substrate in the presence of 50 mM Tris, pH 7.6, 10 mM MgCl₂, 5 mM dithiothreitol, 1-2 mM ATP. If kinasing is for labeling or probe, the ATP will contain high specific activity gamma³²P.

Ligations are performed in 15-30 μ l volumes under the following standard conditions and temperature: 20 mM Tris-Cl pH 7.5, 10 mM MgCl₂, 10 mM dTT, 33 μ g/ml BSA, 10 mM-50 mM NaCl, and either 40 μ M ATP, 0.01-0.02 (Weiss) units T4 DNA ligase at 0°C (for "sticky end" ligation) or 1 mM ATP, 0.3-0.6 (Weiss) units T4 DNA ligase at 14°C (for "blunt end" ligation). Intermolecular "sticky end" ligations are usually performed at 33-100 μ g/ml total DNA concentrations (5-100 nM total end concentration). Intermolecular blunt end ligations (usually employing a 10-30 fold molar excess of linkers) are performed at 1 μ M total ends concentration.

In vector construction employing "vector fragments", the vector fragment is commonly treated with bacterial alkaline phosphatase (BAP) in order to remove the 5' phosphate and prevent religation of the vector. BAP digestions are conducted at pH 8 in approximately 150 mM Tris, in the presence of Na^+ and Mg^{+2} using about 1 unit of BAP per μg of vector at 60°C for about one hour. In order to recover the nucleic acid fragments, the preparation is extracted with phenol/chloroform and ethanol precipitated. Alternatively, religation can be prevented in vectors which have been double digested by additional restriction enzyme digestion of the unwanted fragments.

C.4. Modification of DNA Sequences

For portions of vectors derived from cDNA or genomic DNA which require sequence modifications, site specific primer directed mutagenesis is used. This technique is now standard in the art, and is conducted using a primer synthetic oligonucleotide complementary to a single stranded phage DNA to be mutagenized except for limited mismatching, representing the desired mutation. Briefly, the synthetic oligonucleotide is used as a primer to direct synthesis of a strand complementary to the phage, and the resulting double-stranded DNA is transformed into a phage-supporting host bacterium. Cultures of the transformed bacteria are plated in top agar, permitting plaque formation from single cells which harbor the phage.

Theoretically, 50% of the new plaques will contain the phage having, as a single strand, the mutated form; 50% will have the original sequence. The plaques are hybridized with kinased synthetic primer at a temperature which permits hybridization of an exact match, but at which the mismatches with the original strand are sufficient to prevent hybridization. Plaques which hybridize with the probe are then picked, cultured, and the DNA recovered. Details of site specific mutation procedures are described below in specific examples.

C.5. Verification of Construction

In the constructions set forth below, correct ligations for plasmid construction are confirmed by first transforming E. coli strain MM294, or other suitable host with the ligation mixture. Miniprep DNA was prepared according to Ish-Horowicz, D. et al. (Nucl. Acids Res. (1981) 9:2989) and screened by restriction analysis. DNA may be further analyzed by restriction and/or sequenced by the dideoxy method of Sanger, F., et al., Proc. Natl. Acad. Sci. (USA) (1977) 74:5463 as further described by Messing, et al., Nucleic Acids Res. (1981) 9:309, or by the method of Maxam, et al., Methods in Enzymology (1980) 65:499.

C.6. Transfer Vector Construction

Because the genome of AcNPV is so large (125kb), there are too many restriction sites to allow site-specific insertion of heterologous genes. Therefore, it is necessary to derive recombinant virus, containing the gene to be expressed, through homologous recombination between viral DNA and genetically engineered chimeric plasmids called transfer vectors.

The transfer vectors which have been described by Smith, G.E., et al., 1983, supra, were originally constructed by cloning the AcNPV EcoRI-1 fragment containing the polyhedrin gene into the EcoRI site of E. coli plasmid pUC8 (Vieira, J., et al., Gene 19:259-268 (1982)). A series of plasmids or transfer vectors having single BamHI cloning sites at various locations in the polyhedrin gene were created as described (Smith et al., 1983, supra). One of these, pAc373, has a single BamHI site 50 bp downstream from the polyhedrin cap site i.e., 8 bp before the polyhedrin ATG translation initiation codon (Smith et al., (1985), supra and in U.S. Patent No. 4,745,051). The transfer vectors, pAc610 and pAc611 have the polylinker from M13mp10 and M13mp11, respectively, inserted at this BamHI site of pAc373 (Luckow, V.A. et al. (1988) Biotechnol. 6:47-55. Partial nucleotide sequence of pAc401 and pAc436 transfer vectors for the production of polyhedrin/foreign gene fusion proteins has been reported (Summers, M.D. et al., 1987, p. 53, supra).

C.7. Preparation and Isolation of Recombinant AcNPV Virus

Detailed methods for the generation of recombinant virus can be found in European Patent Application No. 0127839 to G.E. Smith and M.D. Summers of the Texas A & M University System. In general, 2 μ g of genetically engineered transfer vector DNA and 1 μ g of AcNPV viral DNA are cotransfected onto monolayer culture cells of Spodoptera frugiperda. The infected cells usually show viral occlusions by day 3 or 4, with 10-50% of the cells being infected. The virus titer of the medium is expected to be 10^7 pfu/ml and 0.1%-0.5% are expected to be recombinant virus.

Several methods for the detection of recombinant virus are known in the art. Visual detection of the plaques is best achieved using a low power dissecting microscope and observing the plaques on inverted plates with a black background and illumination from the side. More unequivocal methods for detecting recombinants are plaque hybridization using DNA probes to the cloned gene. Antibody probes to the product of the cloned gene may also be employed.

Isolation of the recombinant virus is achieved through plaque purification of serially infected monolayer cells overlaid with soft agar. After two or three cycles the recombinant virus would be seen as separate plaques showing the characteristic occlusion-negative morphology. The plaques, containing about 10,000 pfu of virus, are picked using a sterile Pasteur pipet and transferred to 2 ml of medium.

C.8. Electrophoretic Analysis of Expression Products

In order to concentrate expression products, culture supernatants were incubated with Protein A-Sepharose CL-4B beads cross-linked to antibody to either CSF-1 or G-CSF for 1 hour. The beads were then pelleted and the supernatants reserved. The beads were washed once in .5 ml buffer containing 1M LiCl, 20 mM Tris-HCl pH 8.0 and 0.5% NP40, three times in .5 ml buffer containing 150 mM NaCl, 20 mM Tris-HCl pH8 and 0.5% NP40 and then suspended in sample loading buffer lacking reducing agent. The samples were then heated to 37°C

for 5 minutes, pelleted and the supernatant removed to new tubes. These supernatants were adjusted to contain 0.14 M beta-mercaptoethanol or 20 mM DTT, heated to 100°C for 4 min. and run on SDS-PAGE. SDS-PAGE was performed essentially according to the procedure of Laemmli (Nature (1970) 227:680-685).

Immunoblotting procedures have been described in commonly owned EP Publication No. 219,286 published April 22, 1987 essentially as described below. Immunoblotting of the gel onto nitrocellulose (Schleicher and Schuell, 0.45 μ meter) was performed in a Bio-rad Trans-blot cell at 35 V for one hour at room temperature essentially according to published methods (Towbin et al. (1979) Proc. Natl. Acad. of Sci., USA, 76:4350-4354; Bittner et al. (1980) Anal. Biochem., 102:459-0471; Burnette et al. (1981) Anal. Biochem., 112:195-203). Following transfer, nonspecific antibody binding sites on the nitrocellulose were blocked by incubation for 30 minutes at room temperature with gentle agitation in 250 ml of 0.1% Tween 20 in phosphate buffered saline (PBS). Then the blot was washed three times with gentle agitation at room temperature for five minutes each in 250 ml volumes of 0.1% nonfat dry milk, 0.1% ovalbumin in PBS, and incubated with gentle agitation for three hours at room temperature in 5 ml of a 1/500 to 1/1000 dilution in the preceding buffer of rabbit antiserum against G-CSF (supplied by BABC0). After washing three times as described above, the blot was incubated for one hour at room temperature with 5 ml of a 1/2000 dilution of goat anti-rabbit IgG conjugated to horseradish peroxidase (this conjugate supplied by Zymed) and washed again three times as above.

In order to concentrate expression products, culture supernatants were incubated with Protein A-Sepharose CL-4B beads cross-linked to antibody to a CSF protein for 1 hour. The beads were then pelleted and the supernatants reserved. The beads were washed once in .5 ml buffer containing 1M LiCl, 20 mM Tris-HCl pH 8.0 and 0.5% NP40, three times in .5 ml buffer containing 150 mM NaCl, 20 mM Tris-HCl pH8 and 0.5% NP40 and then suspended in sample loading buffer lacking reducing agent. The samples were then heated to 37°C for 5 minutes, pelleted and the supernatant removed to new tubes. These

supernatants were adjusted to contain 0.14 M beta-mercaptoethanol or 20 mM DTT, heated to 100°C for 4 min. and run on SDS-PAGE.

C.9. Insect Cell Culture

5 Methods for insect cell cultures are well known in the art and detailed procedures for their cultivation can be found in Summers, M.D. et al. (1987, supra), EP Publication No. 127,839 to G.E. Smith et al. or in U.S. Patent No. 4,745,051 (supra). The insect expression host of the current invention, Spodoptera frugiperda (Sf9) is well suited to the production of heterologous proteins because of its
10 ability to grow in either monolayer or suspension culture.

 As monolayer cultures, Spodoptera frugiperda cells will divide every 18-24 hours depending on the culture media. The cells do not require carbon dioxide to maintain the pH of the medium and they will grow well at temperatures between 25-30°C. Subculturing is done
15 2 or 3 times a week when the cells are confluent. Because insect cells are loosely adherent they are easily resuspended without the need of proteases.

 Suspension culture conditions will vary depending on the medium and culture volume and should be determined empirically.
20 Subculturing is required when the cell density reaches 2×10^6 cells/ml by replacing 80% or more of the culture with an equal volume of fresh medium. With suspension cultures larger than 500 ml it becomes necessary to aerate by either bubbling or diffusion.

 Preferred media and culture conditions have been described
25 by Inlow, D. et al. (1987) "Large-scale insect culture for recombinant protein production, "Presented at Symposium on Strategies in Cell-Culture Scale-Up at the American Chemical Society National Meeting in New Orleans, Louisiana. Appropriate serum-free media include:

(a) a basal medium;

30 (b) a lipid component; and

(c) a peptone component.

A "basal medium" is herein defined as a nutrient mixture of inorganic salts, sugars, amino acids, optionally also containing vitamins, organic acids and/or buffers. Basal media together with supplements provide the nutrients necessary to support cell life, growth and reproduction. The preferred basal media used as the starting point for the preparation of the serum free media of the present invention contain neither serum, nor proteins, nor preferably any peptones. The choice of basal medium for the preparation of the media of this invention is not critical. The basal medium can also be considered optimal in the sense that appropriate peptone and lipid components can be selected which provide such necessary nutrients as amino acids and vitamins required to support cell life, growth and reproduction.

As indicated, IPL-41 is a preferred basal medium for the preparation of the media used in this invention. IPL-41 basal medium is commercially available from a number of vendors and is described in Weiss, et al. In Vitro, 17(6):495-502 (June 1981) and in Weiss et al., CRC Press, supra, pp. 70-72 (1986). Table 1 of Weiss et al. (In Vitro at page 496, and Table 3 of Weiss et al. CRC Press, at pages 71-72 outline all the components of IPL-41 and provide their proportions in mg/l; said tables are herein incorporated by reference. At page 497 of Weiss et al. (In Vitro), the preparation of the complete medium IPL-41 is described wherein tryptose phosphate broth (TPB) and fetal bovine serum (FBS) are added. The IPL-41 basal medium employed in preparing the serum-free media of this invention does not contain tryptose phosphate broth (TPB) or fetal bovine serum (FBS).

The serum-free media of this invention can further comprise a protective agent. Such a protective agent is a preferred component of the media of this invention, especially under well-aerated culture conditions. Therefore, the serum-free media of this invention include:

- (a) a basal medium;
- (b) a lipid component;

(c) a peptone component; and

(d) a protective agent.

5 A protective agent is necessary to prevent cell damage and death under well-aerated conditions as found in well agitated and sparged cultures. The protective agent prevents a disintegration/clumping phenomenon of insect cells growth in shake flasks and adherence of the cells to the vessel walls. Further, the protective agent reduces the amount of cellular debris in shake flask cultures indicating that cell lysis is reduced by the presence of the protectant. The protective agent further preferably acts as an anti-foaming agent preventing the loss of cells from the free suspension into a foam layer, and can act as a bubble surface tension reducing agent and/or as a cell surface stabilizing agent and/or as a viscosifying agent.

15 Protective agents are herein defined as non-toxic, water soluble compounds that functionally act to protect insect cells from damage and death in agitated and sparged insect cell culture. The protective agents of this invention are preferably non-toxic, water soluble polymers. A protective agent candidate can be selected by first confirming that it is not toxic to the insect cells to be cultured by methods known to those skilled in the art of insect cell culture, for example, by adding it to a suspension or monolayer of the insect cells of choice for cultivation and comparing the growth of the culture to a control. Then, the non-toxic protective agent candidates can be tested for protective ability by adding the candidate agent to an agitated or sparged culture of the insect cells of choice at small scale and observing viability and growth rate of the cells of said culture to the viability and growth rate of the cells in a control culture.

30 The protective agents in the media of this invention are preferably cell surface stabilizing agents and/or viscosifying agents and/or bubble surface tension reducing agents. Examples of preferred protective agents are hydroxyethyl starch, methyl cellulose, carboxymethyl cellulose (as, sodium carboxymethyl cellulose),


dextran sulfate, polyvinylpyrrolidone, ficoll, alginic acid, polypropyleneglycol, and non-toxic polymeric detergents. Preferred non-toxic, non-ionic polymeric detergents are block polymers of propylene oxide and ethylene oxide (polyoxypropylene polyoxyethylene condensates), preferably Pluronic polyols, such as, Pluronic F68, F77, F88 and F108, preferably F68 and F88, more preferably F68. The Pluronic polyols are commercially available from BASF Wyandotte Corp. (101 Cherry Hill Road, P.O. Box 181, Parsippany, NJ 07054, U.S.A.).

If the protective agent is not also functioning as an emulsifier, the media of this invention further comprise another emulsifier that acts to emulsify the lipid component in conjunction with another emulsifier present at a small concentration in the lipid component itself as described below. Such an emulsifier alternative to a protective agent/emulsifier is preferably a detergent, preferably non-ionic which is non-toxic to the insect cell culture at the concentrations required for emulsification of the lipid component.

The introduction of the lipid component in the form of a microemulsion enhances the availability of lipids in the media to the cells. One option for emulsifying the lipid component is a dual emulsifier system wherein the protective agent is an emulsifier as well as a protective agent and can act in conjunction with an emulsifier or combination of emulsifiers present in the lipid/organic solvent solution making up the lipid component of the media of this invention. Another option for emulsifying the lipid component of the media of this invention is a system wherein the protective agent is not significantly emulsifying but wherein one or more additional emulsifiers are present in an aqueous solution which is added to the lipid component organic solution and act in conjunction with the emulsifiers present therein to form a microemulsion.

Preferred emulsifiers added to the lipid component of the media of this invention include phospholipids, preferably lecithin and non-toxic non-ionic polymeric detergents, preferably a polysorbate compound having the formula:

28



wherein R is a saturated or unsaturated fatty acid having from 16 to 20 carbons, inclusively

wherein t is an integer between 10 and 30, inclusively; and

wherein u is an integer between 10 and 20, inclusively.

5 Most preferably the non-toxic, non-ionic, polymeric detergent/emulsifier is polyoxyethylene (20) sorbitan monooleate, otherwise known as polysorbate 80. Such a non-toxic, non-ionic polymeric detergent is commercially available as Tween 80 from ICI
 10 Americas Inc. (New Murphy Road & Concord Pike, Wilmington, DE 19897, USA). Another polysorbate 80 is commercially available as Durfax 80 from Durkee Industrial Foods Group/SCM Corp. (900 Union Commerce Bldg., Cleveland, Ohio 44115, USA). Other non-toxic, non-ionic, polymeric detergent candidate emulsifiers can be found in editions of McCutcheon's Emulsifiers and Detergents, supra.

15 Said non-ionic, non-toxic, polymeric detergent/emulsifier, such as, polysorbate 80, is present in the media used in this invention at a concentration from about 5 mg/l to about 75 mg/l, more preferably from about 20 mg/l to about 30 mg/l, and most preferably about 25 mg/l.

20 A preferred example of a dual emulsifier system of the media of this invention is the combination of a protective agent/emulsifier, preferably a Pluronic polyol, more preferably Pluronic F68 or Pluronic F88, and still more preferably Pluronic F68, and a non-toxic, non-ionic polymeric detergent, preferably a polysorbate compound, and more
 25 preferably polysorbate 80.

The lipid component is preferably added to the bulk of the media of the invention as a microemulsion. An advantage of the lipid component being in the form of a microemulsion, in addition to enhancing the availability of the lipids to the insect cells, is in providing the option of not having to filter sterilize the lipid component and the rest of the media components separately. The lipid component in the form of a microemulsion can be added to the media without being filter sterilized, and the entire media can then be filter sterilized without the concern that lipids, for example, in globular form, could be lost during the filter sterilization process. For large scale production, such an advantage is significant.

An exemplary lipid component of the media used in this invention comprises per liter of media:

10 mg cod liver oil

25 mg Tween 80

41.5 mg cholesterol

2 mg alpha-tocopherol

1 ml ethanol.

To the optionally filtered/sterilized lipid component solution (1 ml), then, in this exemplification, 10 ml of 10% Pluronic F68 in water (optionally filtered/sterilized) is slowly added with agitation as by vortexing.

The peptone component of the serum-free media of this invention can be selected from a wide variety of hydrolyzed protein products, either alone or in combination, including, without limitation, ox liver digest, such as Panmede (commercially available from Paines & Byrnes Ltd., Greenford, England), yeast extract, such as Yeastolate (preferably TC Yeastolate from Difco, USA), caseine digest, such as Bactocasitone (commercially available from Difco, USA), tryptose phosphate broth (TPB) wherein tryptose is the peptone, Lactalbumin Hydrolyzate (LH) (commercially available from Difco, USA) gelatin peptone, glycerin-gelatine peptone, and beef peptone among many other proteolytic digest products of proteins.

Preferably the peptone component is composed of peptone fractions either alone or in combination selected from the group comprising TPB, caseine digest preferably Bactocasitone, ox liver digest preferably Panmede, yeast extract, preferably Yeastolate, and Lactalbumin Hydrolyzate (LH). More preferably, the peptone component comprises either LH or Yeastolate, alone or in combination. Still more preferably, the peptone component comprises either a combination of LH and Yeastolate or Yeastolate alone.

The total peptone concentration present in the media of this invention can be as high as the sum of the highest concentrations of the individual peptone fractions wherein said highest concentration for each peptone fraction is that which is non-toxic and non-inhibitory to cell growth and wherein the total peptone concentration of said highest, non-toxic, non-inhibitory concentrations of the peptone fractions is similiary non-toxic and non-inhibitory to cell growth. The highest peptone concentrations can vary not only with the particular peptone fractions used but also with the particular insect cell line that is selected. In general, preferred total peptone concentration present in the media of this invention can range from about 1 g/l to about 12 g/l, more preferably from about 2 g/l to about 8 g/l, and still more preferably from about 3 g/l to about 5 g/l.

Example I

A. Construction of New Baculovirus Transfer Vectors

A.1. Construction of pAcC1

pAcC1 is similar to pAc401 (described previously in Section C.5. except that the recognition site for EcoRI endonuclease has been removed. To accomplish this, pAc401 was digested to completion with EcoRI and the ends were made blunt using Klenow fragment. After ligation and transformation, candidates were screened for the absence of an EcoRI site.

A.2. Construction of pAcC2

pAcC2 is similar to pAc436 (described previously in Section C.5. except that the recognition site for EcoRI endonuclease has been removed. To accomplish this, pAc436 was digested to completion with EcoRI and the ends were made blunt using Klenow fragment. After ligation and transformation, candidates were screened for the absence of an EcoRI site.

A.3. Construction of pAcC3

pAcC3 differs from pAcC2 in that an NcoI restriction site has been introduced at the ATG translational start of the polyhedrin gene. To accomplish this the new transfer vector, pAcC2, was digested to completion with SmaI endonuclease. Following phenol extracton and ethanol precipitation, SmaI digested pAcC2 was dissolved in TE buffer (10 mM Tris-HCl pH 7.4; 1 mM EDTA). In a 50 μ l volume of ExoIII buffer (50 mM Tris-HCl pH 8.0; 5 mM MgCl₂; 10 mM beta-mercaptoethanol), 10 μ g of SmaI digested pAcC2 was treated with 50 units of E. coli Exonuclease III (ExoIII) at 30°C for 5 minutes. The sample was phenol extracted and ethanol precipitated twice. Then 50 pmoles of a primer EK85, 5'AACCTATAAACCATGGCGGCCCGG3', was kinased with cold ATP in a 20 μ l reaction volume (50 mM Tris-HCl pH 7.8; 10 mM MgCl₂; 10 mM beta-mercaptoethanol). To 5 μ g of ExoIII treated pAcC2 was added 10 pmoles of kinased EK85 in a final volume of 20 μ l NET (100 mM NaCl; 10 mM Tris-HCl pH 7.5; 1 mM EDTA) buffer. To anneal the plasmid and primer, the reaction was heated to 65°C for 10 minutes, incubated at 37°C for 10 minutes and placed on ice. The extension reaction was performed by adding 20 μ l 2 x Klenow buffer (40 mM Tris-HCl pH 7.5; 20 mM MgCl₂; 2 mM beta-mercaptoethanol) containing 1 μ l 10 mM dNTPs, 1 μ l 10 mM ATP, 1 μ l (about 2 units) Klenow fragment and 1 μ l (about 1-2 units) T4 DNA ligase. The reaction was incubated at 16°C for about 4 hours and then transformed into MM294. Minilysates were screened by analyzing for the presence of an NcoI site. Miniprep DNA was then used to retransform and obtain the desired pure clone.

A.4 Construction of pAcC4 and pAcC5

pAcC4 and pAcC5 are derivatives of pAcC3 containing a polylinker sequence at the SmaI site. The polylinker contains recognition sites for restriction endonucleases SmaI, KpnI, PstI, BglII, XbaI (cleavable when DNA is unmethylated), EcoRI, BamHI and BclI. pAcC4 contains the sequence in one orientation while pAcC5 contains the polylinker in the opposite orientation (see Figure 4). To construct these vectors pAcC3 was digested to completion with XmaI endonuclease and ligated with two complementary self-annealed oligomers having the sequence

5'-CCGGGTACCTGCAGATCTAGAATTCGGATCCTGATCA-3'
3'-CATGGACGTCTAGATCTTAAGCCTAGGACTAGTGGCC-5'

After transformation of MM294, miniprep DNAs of transformants were analyzed for the presence of restriction sites in the polylinker sequence.

Example II

A. Isolation of cDNA Encoding Human CSF-1

A.1 pcCSF-17

The human derived pancreatic carcinoma cell line MIA PaCa-2 was used as a source of mRNA to validate probes and for the formation of a cDNA library containing an intronless form of the human CSF-1 coding sequence. The MIA PaCa cell line produces CSF-1 at a level approximately 10 fold below that of the murine L-929 cells.

Negative control mRNA was prepared from MIA PaCa cells maintained in serum-free medium, i.e. under conditions wherein they do not produce CSF-1. Cells producing CSF-1 were obtained by reinducing CSF-1 production after removal of the serum.

Cells were grown to confluence in roller bottles using Dulbecco's Modified Eagles' Medium (DMEM) containing 10% fetal calf serum, and produced CSF-1 at 2000-6000 units/ml. The cell cultures were washed, and reincubated in serum-free DMEM to suppress CSF-1 formation for negative controls. No detectable CSF-1 was produced

after a day or two. Reinduced cells were obtained by addition of phorbol myristic acetate (100 ng/ml) to obtain production after several days of 1000-2000 units/ml.

5 The mRNA was isolated by lysis of the cell in isotonic buffer with 0.5% NP-40 in the presence of ribonucleoside vanadyl complex (Berger, S. L., et al. Biochemistry (1979) 18:5143) followed by phenol chloroform extraction, ethanol precipitation, and oligo dT chromatography, and an enriched mRNA preparation obtained. In more detail, cells were washed twice in PBS (phosphate buffered saline) and are resuspended in IHB (140 mM NaCl, 10 mM Tris, 1.5 mM MgCl₂, pH 8) containing 10 mM vanadyl adenosine complex (Berger, S. L. et al. supra).

15 A non-ionic detergent of the ethylene oxide polymer type (NP-40) was added to 0.5% to lyse the cellular, but not nuclear membranes. Nuclei were removed by centrifugation at 1,000 x g for 10 minutes. The postnuclear supernatant was added to two volumes of TE (10 mM Tris, 1 mM ethylenediaminetetraacetic acid (EDTA), pH 7.5) saturated phenol chloroform (1:1) and adjusted to 0.5% sodium dodecyl sulfate (SDS) and 10 mM EDTA. The supernatant was reextracted 4 times and phase separated by centrifugation at 2,000 x g for 10 minutes. The RNA was precipitated by adjusting the sample to 0.25 M NaCl, adding 2 volumes of 100% ethanol and storing at -20°C. The RNA was pelleted at 5,000 x g for 30 minutes, washed with 70% and 100% ethanol, and was then dried. Polyadenylated (poly A⁺) messenger RNA (mRNA) was obtained from the total cytoplasmic RNA by chromatography on oligo dT cellulose (Aviv, J., et al., Proc. Natl. Acad. Sci. (1972) 69:1408-1412). The RNA was dissolved in ETS (10 mM Tris, 1 mM EDTA, 0.5% SDS, pH 7.5) at a concentration of 2 mg/ml. This solution was heated to 65°C for 5 minutes, then quickly chilled to 4°C. After bringing the RNA solution to room temperature, it was adjusted to 0.4 M NaCl and was slowly passed through an oligo dT cellulose column previously equilibrated with binding buffer (500 mM NaCl, 10 mM Tris, 1 mM EDTA, pH 7.5 0.05% SDS). The flowthrough was passed over the column twice more. The column was then washed with 10 volumes of binding buffer. Poly A⁺ mRNA was eluted with aliquots of ETS, 35

extracted once with TE-saturated phenol chloroform and was precipitated by the addition of NaCl to 0.2 M and 2 volumes of 100% ethanol. The RNA was reprecipitated twice, was washed once in 70% and then in 100 % ethanol prior to drying.

5 Total mRNA was subjected to 5-20% by weight sucrose gradient centrifugation in 10 mM Tris HCl, pH 7.4, 1 mM EDTA, and 0.5% SDS using a Beckman SW40 rotor at 20°C and 27,000 rpm for 17 hours. The mRNA fractions were then recovered from the gradient by ethanol precipitation, and injected into Xenopus oocytes in the standard translation assay. The oocyte products of the RNA fractions were assayed in the bone marrow proliferation assay (or in the bone marrow proliferation assays of Moore, R. N., et al., J. Immunol. (1983) 10 131:2374, and of Prystowsky, M. B., et al., Am. J. Pathol. (1984) 114:149) and the fractions themselves were assayed by dot blot hybridization to a 32-mer probe corresponding to the DNA in the second exon of the genomic sequence (exon II probe). The overlining in 15 Figure 1 shows the exon II probe.

 The bone marrow proliferation assay results of the supernatants from the Xenopus oocytes did not correlate exactly with 20 the dot-blot results. The most strongly hybridizing fraction, 11, corresponds to 18S, while the most active fractions 8 and 9 correspond to 14-16S. Fraction 8, 9 and 11 were used to form an enriched cDNA library as described below.

 The mRNA was also fractionated on a denaturing formaldehyde gel, transferred to nitrocellulose, and probed with exon II probe. 25 Several distinct species ranging in size from 1.5 kb to 4.5 kb were found, even under stringent hybridization conditions. To eliminate the possibility of multiple genes encoding CSF-1, digests of genomic DNA with various restriction enzymes were subject to Southern blot and 30 probed using pcCSF-17 DNA. The restriction pattern was consistent with the presence on only one gene encoding CSF-1.)

 The enriched mRNA pool was prepared by combining the mRNA from the gradient fractions having the highest bone marrow proliferative activity, although their ability to hybridize to probe

is relatively low (14S-16S) with the fractions hybridizing most intensely to probe (18S). Higher molecular weight fractions which also hybridized to exon II probe were not included because corresponding mRNA from uninduced MIAPaCa cells also hybridized to exon II probe.

5 cDNA libraries were prepared from total or enriched human mRNA in two ways. One method uses gt10 phage vectors and is described by Huynh, T. V., et al., in DNA Cloning Techniques: A Practical Approach IRL Press, Oxford 1984, D. Glover, ed.

10 A preferred method uses oligo dT priming of the poly A tails and AMV reverse transcriptase employing the method of Okayama, H., et al., Mol. Cell Biol. 1(1983) 3:280-289, incorporated herein by reference. This method results in a higher proportion of full length clones than does poly dG tailing and effectively uses as host vector portions of two vectors therein described, and readily obtainable from
15 the authors, pcDV1 and pL1. The resulting vectors contain the insert between vector fragments containing proximal BamHI and XhoI restriction sites; the vector contains the pBR322 origin of replication, and Amp resistance gene and SV40 control elements which result in the ability of the vector to effect expression of the
20 inserted sequences in COS-7 cells.

A 300,000 clone library obtained from above enriched MIAPaCa mRNA by the Okayama and Berg method was then probed under conditions of high stringency, using the exon II probe. Ten colonies hybridizing to the probe were picked and colony purified. These clones were
25 assayed for the presence of CSF-1 encoding sequences by transient expression in COS-7 cells. The cloning vector, which contains the SV40 promoter was used per se in the transformation of COS-7 cells.

30 Plasmid DNA was purified from the 10 positive clones using a CsCl gradient, and the COS-7 cells transfected using a modification (Wang, A. M., et al., Science (1985) 228:149) of the calcium phosphate coprecipitation technique. After incubation for three days, CSF-1 production was assayed by subjecting the culture supernatants to the radioreceptor assay performed substantially as disclosed by Das, S. K., et al., Blood (1981) 58:630, and to a colony stimulation (bone

marrow proliferation) assay performed substantially as disclosed by Prystowsky, M. B., et al., Am. J. Pathol. (1984) 114:149. Nine of the ten clones picked failed to show transient CSF-1 production in COS-7 cells. One clone, which did show expression, was cultured, the plasmid DNA isolated, and the insert was sequenced. The DNA sequence, along with the deduced amino acid sequence, are shown in Figure 1. The full length cDNA is 1.64 kb and encodes a mature CSF-1 protein of 224 amino acids. The clone was designated CSF-17 with Cetus depository number CMCC 2347 and was deposited with the American Type Culture Collection on June 14, 1985, as accession No. 53149. The plasmid bearing the CSF-1 encoding DNA was designated pcCSF-17.

A.2. Additional Clones

pcCSF-17, prepared as described above, was used as a probe to obtain additional CSF-1 encoding clones from a human cDNA library. Total mRNA was prepared from MIAPaCa cells exactly as described above and used to obtain a cDNA library in gt10 phage vectors by ligating the reverse transcripts to EcoRI linkers and inserting the EcoRI digest of the cDNA thus provided into the EcoRI site of ggt10, as described by Huynh, T. V., et al., in DNA Cloning Techniques: A Practical Approach, IRL Press, Oxford, 1984, D. Glover, ed.

Over one million phage were screened using a single-stranded highly labeled probe derived from CSF-17 using standard procedures, which are briefly summarized as follows.

A BglI-AvaII fragment of pcCSF-17 DNA, which includes the entire coding sequence for CSF-1 plus about 600 bp of 3' untranslated region was inserted into M13, and a labeled second strand synthesized as follows: Approximately 1 pmol of M13 containing the single-stranded BglI-AvaII-digested pcCSF-17 was annealed with 10 pmol of sequencing primer in 20 mM Tris, pH 7.9, 10 mM MgCl₂, 100 mM NaCl, and 20 mM beta-mercaptoethanol at 67°C for 5 minutes and then transferred to 42°C for 45 minutes. The annealed preparation was supplied with 100 μ mol each of dATP, dCTP, and dTTP, and 2.5 μ mol P³²-labeled (alpha)GTP, along with 5 U Klenow fragment. The reaction was

incubated at 37°C for 20 minutes, and the DNA recovered on a P-6 dG (Bio-Rad) spin column and boiled for 10 minutes to separate the strands.

The probes, thus prepared, were used to screen the plaques (which had been transferred to nitrocellulose) by hybridization under stringent conditions (50% formamide, 5 x SSC, 5 x Denhardt's) at 42°C for 18 hours.

Approximately 150 phage plaques were positive, and 20 which were particularly strongly hybridized were selected for plaque purification to obtain individual clones.

The 20 clones were then subjected to hybridization under the same conditions to the oligonucleotide GM11, which has the sequence complementary to nucleotides 506-545 in Figure 2. As described above, this sequence is an exact match to that portion of the human genomic sequence which corresponds to the "extra" portion of the murine cDNA, described below, that encodes the "extra" 295 amino acid segment in the "long form" of the murine CSF-1 protein.

The complete DNA sequence for the pertinent coding regions of the cDNA insert in clone 4 along with the deduced amino acid sequence, are shown in Figure 2. The sequence was derived by integrating the known sequence of the genomic clone, phCSF-1a, described above, using the 295 amino acid "extra" insert of the murine sequences described below as a guide to deduce the complete sequence shown. The sequence depicted shows the splicing of the "extra" segment, sufficient to encode 298 "extra" amino acids contained in the gene at the 5' side of exon 6, into the sequence of pcCSF-17 between the first nucleotide of the codon for the Gly residue at amino acid position 150 into reading frame with the remaining CSF-17 sequence. The insert changes the codon at 150 to a codon for aspartic acid, the subsequent codon at the end of the insert is reconstituted to a Gly, and the remaining sequence of residues continuing with His-Glu-Arg etc. down to the C-terminal Val residue remain the same as in pcCSF-17.

Two of the clones, 4 and 25, were cloned into M13mpl8 or M13mpl9 for sequencing to confirm the results shown in Figure 2 (only clone 4 sequence shown) using the EcoRI restriction sites. These two clones appeared identical from restriction enzyme analysis. They were then subclones into the modified Okayama-Berg vector pcDB, which was prepared from the published Okayama-Berg vectors pcDV1 and pL1 (Okayama, H., et al., Mol. Cell Biol. (1983) 3:280-289) as follows:

pcDV1 was digested with HindIII and KpnI and the large fragment, providing Amp^r , the pBR322 origin of replication, and a polyadenylation site, was isolated. pL1 was digested with HindIII and PstI and the small fragment, providing the SV40 promoter and origin of replication, was isolated. These fragments were religated in a three-way ligation with the synthetic, KpnI/PstI-digested oligonucleotide fragment

```
CTGCAGGAGCTCAGATCTTCTAGAGAATTCTCGAGCGGCCGCATCGATGGTACC
GACGTCCTCGAGTCTAGAAGATCTCTTAAGAGCTCGCCGGCGTAGCTACCATGG
```

to obtain pcDB, a plasmid corresponding to pcD-x shown in the reference, wherein "x" is replaced by a polylinker. Thus, pcDB contains the SV40 early promoter and an immediately downstream linker followed by a polyadenylation site. Sequences ligated into the polylinker region should be expressed under the control of the SV40 early promoter.

Before testing expression, because clones 4 and 25 appeared to be missing some 5' end sequences as compared to CSF-17, the upstream portions from pcCSF-17 were substituted for those of clones 4 and 25.

The protocol for this substitution was as follows: pcCSF-17 was digested with SmaI, which cuts at the extreme 5' end of the untranslated sequence (see Figure 1) and the linearized plasmid was ligated to EcoRI linkers and resealed. The religated plasmid was then digested with EcoRI, which removes the entire coding region from just upstream of the SmaI site at the extreme 5' end to the EcoRI site immediately downstream of the stop codon. This was ligated into the polylinker of pcDB at the EcoRI site. The coding sequences downstream

of the BstXI site, which site is located at approximately the codon for amino acid 8 of the CSF-17 mature protein sequence (see Figure 1) was removed by digestion with BstXI and KpnI. (KpnI cuts into the linker slightly downstream of the EcoRI site past the stop codon.) This deleted downstream sequence was replaced by the analogous BstI/KpnI fragment from mp13-4 and mp13-25. The resulting clones, pcDBhuCSF-4 and pcDBhuCSF-25 thus contained the coding sequences downstream of the codon for approximately amino acid 8 from clones 4 and 25, respectively, and the upstream sequences from pCSF-17. The ligated sequences are in reading frame and under control of the SV40 early promoter.

B. Construction of Recombinant Baculovirus Transfer Vectors Containing the CSF-1 Gene

B.1. Construction of pAcM1 and pAcM2

pAcM1 was constructed by taking advantage of the SmaI and EcoRI sites located outside the CSF-1 coding sequence but within the insert containing the CSF-1 gene in pcCSF-17 (see Kawasaki, E. S. et al., (1985) supra; SmaI recognition site begins at bp 13 and EcoRI begins at bp 953). The pAc611 transfer vector was doubly digested with EcoRI and SmaI endonucleases. The 937 bp SmaI-EcoRI fragment from pcCSF-17 containing the CSF-1 coding sequence was isolated by gel electrophoresis and ligated to SmaI-EcoRI digested pAc611 using T4 DNA ligase. E. coli MM294 was transformed with the ligation mixture and transformants were screened for the presence of the 937 bp SmaI-EcoRI CSF-1 fragment.

pAcM2 was constructed in a manner analogous to that described for pAcM1 except that the SmaI-EcoRI fragment from pCSF-BamBcl described in commonly owned PCT Publication No. W086/04607, supra was inserted into pAc611 rather than the SmaI-EcoRI fragment from pcCSF-17. pCSF-BamBcl differs from pcCSF-17 in that a mutation has been introduced to change the serine at position 159 in the CSF-1 coding sequence to a stop codon. This mutation was accomplished by excising the coding sequence from pcCSF-17 and ligating into M13 for site-specific mutagenesis using the primer,

5'-GAGGGATCCTGATCACCGCAGCTCC-3'.

This results in a new BclI site at codons 159-160. The mutated DNA was excised with BstXI and EcoRI and ligated into BstXI-EcoRI digested pcCSF-17. The ligation mixture was transformed into E. coli DG105, a dam⁻ host, and the plasmid DNA isolated. This host was used because BclI is sensitive to dam methylation. The resulting plasmid, pCSF-BamBcl, was the source of the CSF-1 sequence present in pAcM2.

B.2. Construction of pAcM3

pAcM3 was constructed by using M13 site-specific oligonucleotide directed mutagenesis of the pAcM2 vector to delete the CSF-1 5' untranslated leader sequence and to reconstruct the AcNPV polyhedrin gene 5' untranslated leader sequence. The approximately 900 bp EcoRV-EcoRI fragment from pAcM2 was ligated into SmaI-EcoRI digested M13RF. The EcoRV site is located at -96 bp in the polyhedrin 5' leader (see Smith, G. E. et al., Mol. Cell. Biol. 3:2158) and the EcoRI site begins at bp953 following the CSF-1 coding sequence as previously described in Section B1. Using the primer,

5'-GTTTTGTAATAAAAAACCTATAAATAATGACCGCGCCGGGC-3'

the polyhedrin 5' untranslated leader is juxtaposed with the ATG translational start of CSF-1. The CSF-1 5' untranslated leader sequence as well as the polylinker sequence of the transfer vector are deleted. The mutated DNA sequence (EK82RF) was restricted with XbaI, the ends were made blunt by Klenow repair and a second endonuclease digestion with EcoRI resulted in a XbaI(blunt)-EcoRI fragment which was inserted into EcoRV-EcoRI digested pAc611. After ligation and transformation into DH5, transformants were screened by restriction analysis. The desired construct designated pAcM3 contains a small insertion in the polyhedrin promoter region at the EcoRV site as a result of subcloning the M13 fragment back into pAc611.

B.3. Construction of pAcM4

pAcM4 is an in frame translational fusion vector constructed by cloning a mutated CSF-1 cDNA fragment into pAcC1. The source of

the CSF-1 cDNA fragment was pCSF-gly150 described in PCT Publication No. W086/04607, supra which contains a TGA stop codon instead of histidine codon at position 151. It was prepared from the pcCSF-17 insert by site-specific mutagenesis using the appropriate primer,

5'-AGCCAAGGCTGATCAAGGCAGTCC-3'.

Thus, the downstream portion of the coding sequence was excised from pcCSF-17 with BstXI-EcoRI, put into an M13 vector for mutagenesis and returned after plaque purification as a BstXI-EcoRI insert. The resulting plasmid is designated pCSF-gly150.

The SmaI-EcoRI fragment (bp 13-953) in pCSF-gly150 was then inserted into an M13 vector for a second round of mutagenesis. The objective this time was to introduce a SmaI recognition site starting 3 bp after the ATG translational start codon of the CSF-gly150 coding sequence. This was accomplished by site-specific mutagenesis using the appropriate primer 5'-GCCCGTATGTCCCCGGGGGCGCCG-3'. The mutagenized DNA in M13 is designated EK83RF.

EK83RF was used as the source of the CSF mutein used in the construction of pAcM4. EK83RF was digested with EcoRI endonuclease, repaired with Klenow fragment and digested with SmaI endonuclease. The 770 bp fragment containing the CSF sequence was gel purified and ligated into SmaI digested pAcC1, using T4 DNA ligase. The ligation mixture was used to transform MM294 and transformants were screened by minilysates or colony hybridization.

The desired construct pAcM4 thus contains the intact polyhedrin gene promoter and 5' untranslated leader but three altered codons in the CSF-1 signal peptide. Instead of Met-Thr-Ala-Pro-Gly...the sequence in this construct encodes Met-Arg-Pro-Gly-Gly... The CSF-1 5' untranslated leader sequence has been removed so that all of the sequence 5' to the ATG translational start is derived from the AcNPV polyhedrin gene (see Figure 5).

B.4 Construction of pAcM5 and pAcM6 Containing "Long Form" CSF-1

pAcM5 was prepared by placing the coding region of CSF-4 from pcDBhuCSF-4 into pAc610. pcDBhuCSF-4 was digested with EcoRI

and the 1826 bp fragment containing the 522 amino acid coding sequence of the "long form" of CSF-1 was isolated by gel electrophoresis. The pAc610 transfer vector was also digested with EcoRI endonuclease and ligated with the EcoRI fragment containing the CSF-1 gene. E. coli MM294 were transformed with the ligation mixture and transformants were screened for hybridization to the oligonucleotide. GM11 (described in Section A.2) which hybridizes to the extra amino acid sequence not found in pcCSF-17. The resulting construct was designated pAcM5.

pAcM6 was constructed by excising the 250 bp StuI (positions 511-821 bp on the pcCSF-17 sequence) fragment from pAcM4 (described in Section B.3) and replacing it with the 1141 bp StuI (positions 343-1484 bp on the CSF-4 sequence) fragment from pcDBhuCSF-4. The 1141 bp StuI fragment from pcDBhuCSF-4 contains the sequence from within amino acid 101 to within amino acid 184 of the shorter pcCSF-17 CSF sequence, with an additional coding sequence for 298 amino acids inserted after amino acid 149. Said in another way, the resulting construction, pAcM6, has the same 5' and 3' sequence as found in pAcM4 with the additional 894 bp of additional coding sequence from pcDBhuCSF-4 inserted within the CSF-1 coding sequence.

B.5. Construction of pAcM9

pAcM9 was constructed by replacing a fragment of pAcM4 with a mutated fragment of CSF-1 cDNA derived from M13 clone EK90/91. The mutations in this fragment alter the coding sequence of pAcM4 such that amino acids Asn122 and Asn140 are replaced with Gln.

EK90/91 was constructed as follows. The 983 bp SmaI-EcoRI fragment from pcCSF-17-asp59-gly150 was cloned into M13mp18 and subjected to oligonucleotide directed mutagenesis using the primer 5'-GAATGTCTTCCAAGAAACAAAGA-3' to produce M13 clone EK90 (codon 122 altered to Gln). EK90 was then mutated with the primer 5'-CAAGAACTGTCAAAACAGCTTTG-3' to produce the double mutant EK90/91 (codon 122 and 140 altered to Gln). The mutated 940 bp SmaI-EcoRI fragment from EK90/91 was then subcloned into pUC18 to product pUC90/91.

The construction pAcM9 was completed by excising the 250bp StuI fragment (511-821bp in pCSF17) from pACM4 and ligating it to the 250bp from pUC90/91. The ligated DNA was transformed into MM294 and the pAcM9 clone was identified by colony hybridization.

5 B.6. Construction of pAcM10

pAcM10 was constructed by inserting a CSF-1 sequence modified by the introduction of an NcoI site at the ATG translational start into the pAcC4 transfer vector. The modified CSF-1 sequence was derived as follows. The double mutant, pCSF-17-asp59-gly150, described in PCT Publication No. W086/04607, supra was constructed by a first site specific mutagenesis to change amino acid 59 to aspartic acid and create a new EcoRV site at codons 59-60 using the primer 5'-GGTACAAGATATCATGGAG-3'. The resulting plasmid, pCSF-17-asp59, was digested with BstXI and EcoRI to place the C-terminal fragment into M13 for a second site-specific mutagenesis with the same primer described in Section B.3 for the generation of pCSF-17-gly150. The mutated fragment was then returned to the vector to obtain pCSF-17-asp59-gly150.

pCSF-17-asp59-gly150 was digested with XmaI (bp 13-18 in pcCSF-17) and NarI (bp 47-52, bp 99-104, bp 191-196 in pcCSF-17) to completion. NarI cut twice within the 5' untranslated sequence of CSF-1 and the large NarI-XmaI fragment containing the CSF-1-asp59-gly150 sequence from amino acid 6 on through the vector was purified by gel electrophoresis. Complementary primers having the sequence

25 5'-CCGGGACCATGGCCGCGCCGGG-3'
3'-CTGGTACCGGCGCGGCCCGC-5'

were self annealed and ligated to the NarI-XmaI fragment containing CSF-1 sequence. Transformants of E. coli MM294 were screened for the presence of the NcoI site introduced in the primer sequence.

30 pAcM10 was constructed by digesting pCSF-17-NcoI with NcoI and BamHI. pAcC4 was also doubly digested with NcoI and BamHI and after fragment isolation, ligation and transformation, minilysates of E. coli DH5 candidates were screened for the appropriate NcoI-BamHI fragment sizes.

B.7. Construction of pAcM11

pAcM11 was produced by replacing a fragment of pAcM6 with a mutated fragment of CSF-1 cDNA derived from pUC90/91 (described above). The mutations in this fragment alter the 522 amino acid coding sequence of pAcM6 such that amino acids Asn122 and Asn140 are replaced with Gln. pUC90/91 and pcDBhuCSF4 (described in Section A.2) were cleaved with both EcoRI and BsmI. The 1137 bp BsmI (position 477) to EcoRI (position 1614) fragment from pcDBhuCSF4 was agarose-gel purified and ligated to the large fragment of pUC90/91. The ligation reaction was transformed into MM294 cells and colonies were screened by hybridization to the oligomer GM11 (described in Section A.2). The resultant plasmid, pUC90/91EB, contains the entire coding region of CSF4 with the changes previously mentioned. The 1138bp StuI fragment from pUC90/91EB (343-1481bp) was exchanged with the corresponding fragment in pAcM6 to produce pAcM11. pAcM11 was transformed into MM294 and isolated by colony hybridization.

B.8. Construction of pAcM12 and pAcM13

pAcM12 and pAcM13 were derived from pAcM6 and pAcM11 respectively by replacing a 721 bp BamHI fragment (between positions 683 and 1404) with the corresponding fragment derived from M13 clone 4/29/33, an m13mp18 mutant of CSF4. The 4/29/33 fragment has a stop codon immediately following the codon for the proline at amino acid position 221. Therefore, pAcM12 encodes a "long form" CSF-1 protein that is truncated after proline 221 "long form" CSF-1 protein that is truncated after proline 221 that may be glycosylated at Asn 122 and Asn 140. pAcM13 encodes a truncated "long form" CSF-1 protein that cannot be glycosylated at those positions. Clone 4/29/33 was produced by oligonucleotide directed mutagenesis of 4/29 with the primer GM33 which changed the Asn122 and Asn140 to Gln residues.

C. Expression of CSF-1 in Insect Cells

Transfection of insect cells, plaque purification of recombinant virus and infection of insect cells (10^5 cells/ml) with the transfer vectors containing CSF-1 described hereinabove, yielded

between 7500 and 500,000 U/ml of CSF-1 activity in small scale infections of 2-25 ml. Expression levels for each of the vectors are compared in Figure 8.

Protein products obtained from the Baculovirus expression system were analyzed by immunoprecipitation of supernatants from infected Sf9 cells. Samples were eluted from rabbit anti-CV1-CSF-1 covalently coupled to Sepharose beads, run on SDS-PAGE and stained with Commasie blue. Figure 6 shows CSF-1 produced by cells cotransfected with pAcM4 (lane 3), pAcM3(lane 4 and pAcM6 (lane 5).

Since the expression products in most cases were precipitated from cell supernatants it is clear that the products were secreted. Molecular weights of the CSF proteins produced indicated that the signal peptide was cleaved though the precise site of cleavage has not yet been determined. The products also appear to be glycosylated as indicated by their increased molecular weight as compared to the CSF produced intracellularly (in the absence of a signal peptide) in E. coli.

Example III

Isolation and Expression of cDNA Encoding Human G-CSF

A cDNA clone encoding human granulocyte stimulating factor was isolated from the MIA PaCa-2 cell line, and was expressed using a recombinant Baculovirus vector in insect cells. The MIA PaCa-2 cell line described in Yunis, A. A. et al., Experimental Hematol., 12:838-843 (1984), is an established cell line publicly available from the Cell Repository Lines (CRL) collection of the American Type Culture Collection, 12301 Parklawn Avenue, Bethesda, MD 20895 under accession number ATCC CRL 1420. The cDNA clone was sequenced and the corresponding amino acid sequence was deduced.

A. Isolation of G-CSF mRNA From MIA PaCa-2 Cells

Confluent MIA PaCa-2 cells were stimulated in serum free Dulbecco's minimum essential medium (DMEM) for 4 days with phorbol

myristate acetate (50 ng/ml) and retinoic acid ($10 \mu\text{M}$). The RNA was prepared as described by Chirguin et al.; briefly, the cells were lysed in 5 M guanidine isothiocyanate followed by centrifugation through a 5.7 M cesium chloride (CsCl) cushion; poly A⁺ RNA was prepared from the lysed cells by one selection cycle on oligo (dT) cellulose as described in Maniatis et al., supra.

In more detail, cells were lysed in a solution which contained 5 M guanidine isothiocyanate, 0.025 M Na-citrate, pH 7, 0.5% sarcosine and 8% beta-mercaptoethanol. Molecular biology grade CsCl was made up to 5.7 M or 40% w/v and buffered with 0.02 M Tris pH 7.5 and 0.002 M Na-EDTA. All solutions were prepared under RNase free conditions and passed through 0.45 Millipore filters before use. The lysed cells were then layered onto SW28 ultracentrifuge tubes which contained layers of 10 ml 5.7 M CsCl and 6 ml 40% CsCl.

After centrifugation at 26,000 rpm for 18 hours, the RNA pellet was dissolved in H₂O and ethanol precipitated twice. Polyadenylated (Poly A⁺) messenger RNA (mRNA) was obtained by chromatography of the total RNA on oligo (dT) cellulose as described in Maniatis, supra (at page 197).

The RNA was dissolved in sterile H₂O and heated to 65°C for five minutes. An equal volume of a solution containing 0.040 M Tris Cl pH 7.6, 1.0 M NaCl, 0.002 M EDTA and 0.2% SDS was quickly added and the sample was cooled to room temperature. The sample was then loaded on an oligo (dT) column that had been equilibrated with a buffer containing 0.020 M Tris pH 7.6, 0.5 M NaCl, 0.001 M EDTA and 0.1% SDS. The flow through was collected, heated to 65°C, cooled to room temperature and passed over the column once more. The column was then washed with 10 volumes of wash buffer (0.02 Tris, 0.1 M NaCl, 0.001 M EDTA, 0.1% SDS). Poly A⁺ mRNA was eluted with aliquots of 0.01 M Tris pH 7.5, 0.001 M EDTA and ethanol-precipitated twice.

334 μg of MIA PaCa-2 Poly A⁺ mRNA was fractionated on 5-25% by weight sucrose gradient centrifugation in 20 mM Tris HCl, pH 7.5, 1 mM EDTA, and 0.5% sarcosine using a Beckman SW40 rotor at 20°C and 27,800 rpm for 16 hours. The mRNA fractions were collected in 400 μl

fractions and ethanol precipitated twice. Fractions were pooled and resuspended in 15 μ l H₂O.

Northern blots were prepared by the electrophoresis of 5 μ g of poly A⁺ RNA per lane or 1 μ l of each pooled RNA fraction through 1% agarose gels containing 0.5 M formaldehyde followed by blotting onto nitrocellulose filters. After baking the filters for 1.5 hours at 80°C, they were prehybridized (5 x SSC, 10 x Denhardt's solution (0.2% polyvinylpyrrolidone, 0.2% Ficoll, 0.2% BSA), 0.1% SDS, 50 mM sodium phosphate pH 7.0, and 100 μ g/ml tRNA) for one hour at 55°C. The blots were hybridized for 16 hours at 55°C in a similar solution that also contained 10% dextran sulfate and 10⁶ cpm per ml of an oligonucleotide probe labeled with gamma-³²P-ATP and polynucleotide kinase. The blot shown in Figure 3B was washed at 55°C in 3 x SSC, 0.1% SDS. The oligonucleotide probe had the sequence 5'-GTAGGTGGCACACAGCTTCTCTG-3' and was designed based on the sequence of the CHU-2 cDNA clone described by Nagata et al., Nature, 319:415-418 (1986).

The RNA pools that were probe-positive in the above assay were translated in Xenopus laevis oocytes by the injection of 50 nl of RNA into each oocyte as described in Gurdon et al., Nature, 233:177-180 (1971). Supernatants of 10 μ l per oocyte were collected after 40 hours and assayed for CSF activity.

The Xenopus laevis supernatants from each hybridization fraction were assayed for CSF activity in a murine bone marrow cell proliferation assay as described in Moore, et al., J. Immunol., 131:2374-2378 (1983). Briefly, in this assay, 5 x 10⁴ murine bone marrow cells per/well were incubated in 96-well plates (12 x 8) with serially diluted Xenopus laevis oocyte supernatants made from positively hybridizing mRNA fractions. After three days, ³H thymidine (0.5 μ Ci/well) was added, and after six hours the cells were harvested and counted in a liquid scintillation counter.

Peak bone marrow proliferation was found in Xenopus laevis oocyte supernatants made with mRNA fractions that were most strongly positive in Northern blots with the above-described oligonucleotide probe.

B. Identification of G-CSF Clones in a MIA PaCa-2 cDNA Library

A cDNA library was prepared from the enriched MIA PaCa-2 mRNA as described in Kawasaki, et al., Science, 30:291-296 (1985). Briefly, the method used oligo (dT) priming of the poly A⁺ tails and AMV reverse transcriptase employing the method of Okayama, H. et al., Mol. Cell Biol., 3:280-289 (1983), which is incorporated herein by reference. This method results in a higher proportion of full length clones than does poly (dG) tailing and effectively uses as host vector portions of two vectors therein described, and readily obtainable from the authors, pcDV1 and pL1. The resulting vectors contain the insert between vector fragments containing proximal BamHI and XhoI restriction sites; the vector contains the pBR322 origin of replication, and ampicillin resistance gene and SV40 control elements which result in the ability of the vector to effect expression of the inserted sequences in COS-7 cells.

A 1.2×10^6 clone library in E. coli obtained from the above enriched MIA PaCa-2 mRNA by the Okayama and Berg method was then probed using the same oligonucleotide probe that yielded a positive signal on the most active pooled MIA PaCa-2 mRNA fractions. To probe the library, E. coli containing the Okayama-Berg vectors were grown up on nutrient medium. Colonies were lifted onto nitrocellulose filter papers and were lysed. DNA was fixed to the filter by treatment for five minutes with 0.5 mM NaOH, 1.5 M NaCl. Filters were then washed twice for five minutes each time with 1.5 M Tris pH 8, 3 M NaCl and were air dried and baked at 80°C for two hours.

The filters for the screening were prehybridized and hybridized to the gamma³²P labeled probe as described above in Section B of the example, but both prehybridization and hybridization were carried out at 50°C. Plasmids pP12 and pP28 were determined to be probe positive and were further characterized.

C. Sequencing of G-CSF MIA PaCa-2 Plasmids cDNA

pP12 was digested with BamHI and subcloned into a M13mp19 vector and sequenced using the dideoxy chain-termination method.

The DNA sequence and predicted protein sequence of pP12 are shown in Figure 4. The cDNA insert in pP12 is 1510 base pairs long excluding the poly (dG) and poly (dA) tails; it contains 11 more bases of 5' untranslated sequence than the CHU-2 G-CSF clone. The major difference between this clone derived from MIA PaCa-2 and CHU-2 cDNA clones of Nagata et al. is a 9 base pair insertion (GTGATGGAG) in the CHU-2 clone that would encode the amino acid residues Val-Ser-Glu just prior to cys-36 in the MIA PaCa-2 G-CSF as indicated by the arrow in Figure 4. There are two other differences; an A at position 588 in the MIA PaCa-2 clone (G in the CHU-2 clone) is a silent third base change, and a T at position 1237 in the MIA PaCa-2 clone (C in the CHU-2 clone) is in the 3' untranslated region.

2. Construction of Recombinant Baculovirus Transfer Vectors Containing the G-CSF Gene

D.1. Construction of pAcG1 (also known as pJD2)

The Baculovirus G-CSF expression vector, pAcG1, was constructed using the pAc610 transfer vector and the G-CSF coding sequence from pP12. pAc610 was digested with EcoRI endonuclease and the sticky ends were made blunt using the Klenow fragment as described. A second digestion with BamHI endonuclease prepared the transfer vector for receipt of the G-CSF coding sequence immediately downstream from the polyhedrin gene 5' leader sequence. pP12 containing the G-CSF coding sequence was digested with NcoI endonuclease which contains within its recognition sequence the ATG translational initiation codon of the G-CSF gene. Following Klenow repair to generate blunt ends, pP12 was digested with BamHI endonuclease. The NcoI (blunt)-BamHI fragment containing the G-CSF gene was purified by gel electrophoresis. After ligation with T4 ligase and transformation into E. coli MM294 candidates were screened by restriction analysis. Double digestion with EcoRV and BamHI endonucleases resulted in two fragments 1.5 and 8 kb in length.

D.2 Construction of pAcG2

The pAcG1 plasmid lacks the native G-CSF leader sequence but does contain a G-C rich segment from the pAc610 polylinker adjacent to the first ATG of the coding sequence. In addition, this construction has a T at the -3 position from the ATG translational start codon which has been reported to result in low translational efficiency (Kozak, M. Cell, (1986) 44:283). The new transfer vector pAcC3 was designed to eliminate the G-C rich region, restore the complete polyhedrin leader and place an A residue at the -3 position to the translational start.

pAcG2 was constructed by introducing an NcoI-SspI fragment from pP12 containing the G-CSF coding sequence into pAcC3. pP12 was digested with SspI endonuclease which cuts some 40 bp 3' to the translational termination codon of G-CSF to produce blunt ends. A second digestion with NcoI endonuclease results in a 654 bp fragment containing the G-CSF coding sequence (minus leader) which was gel purified. pAcC3 was doubly digested with NcoI and SmaI endonucleases and purified by gel electrophoresis for ligation to the G-CSF fragment. After ligation with T4 ligase and transformation into MM294(?), transformants were screened for presence of insert by digesting with restriction endonucleases. The presence of an NcoI restriction endonuclease recognition site ensured that the translational initiation codon had been regenerated.

E. Expression of G-CSF in Insect Cells

Transfection of insect cells, plaque purification of recombinant virus and infection of insect cells (10^5 cells/ml) with pAcG1 yielded about 90,000 units/ml of G-CSF. Expression levels obtained with pAcG2 after similar procedures were about 2.4×10^6 units/ml of G-CSF. These levels are compared to levels of CSF-1 produced in at least two different constructions in Figure 8.

Protein products obtained from the Baculovirus expression system were analyzed by immunoblotting of supernatants from the infected Sf9 cells. Figure 7 shows the results of this experiment.

Other modifications of the above described embodiments of the invention which are obvious to those of skill in the area of molecular biology and related disciplines are intended to be within the scope of the claims that follow.

Deposits

The materials listed below were deposited with the American Type Culture Collection, Rockville, MD, USA (ATCC). The deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure and the Regulations thereunder (Budapest Treaty). Maintenance of a viable culture is assured for 30 years from date of deposit. The organism will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Applicants and ATCC which assures unrestricted availability upon issuance of the pertinent U.S. patent. Availability of the deposited strain is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

	<u>Recombinant Transfer Vector</u>	<u>CMCC#</u>	<u>ATCC#</u>	<u>Deposit Date</u>
	pAcM3			6/12/87
	pAcM4			6/12/87
	pAcM6			6/12/87
	pAcM10			6/12/87
	pAcG2			

WHAT IS CLAIMED IS:

1. A recombinant baculovirus transfer vector capable of introducing a colony-stimulating factor (CSF) gene or portion thereof into a baculovirus genome said transfer vector comprising a portion of the baculovirus genome, a bacterial replication origin and a DNA sequence encoding a CSF under the transcriptional control of a baculovirus promoter.

2. A recombinant baculovirus transfer vector according to claim 1 wherein said baculovirus promoter is the polyhedrin gene promoter.

3. A recombinant baculovirus transfer vector according to claim 2 wherein said DNA sequence encoding CSF is translationally fused with at least two codons of the polyhedrin gene wherein at least one of said codons is an initiation codon.

4. A recombinant baculovirus transfer vector according to claims 1, 2 or 3 wherein said DNA sequence encoding a CSF is selected from the group of plasmids consisting of pcCSF-17, pcDBhuCSF-4 or pP12.

5. A recombinant baculovirus transfer vector according to claim 4 wherein said DNA sequence encoding CSF-1 from pcCSF-17 contains a translational stop codon replacing either histidine 151 or serine 159.

6. A recombinant baculovirus transfer vector according to claim 5 wherein a polylinker sequence and 5' untranslated leader sequence of CSF-1 are deleted thereby bringing the polyhedrin gene promoter into juxtaposition with the CSF-1 translational initiation codon.

7. A recombinant baculovirus transfer vector according to claim 5 wherein the codons for the second, third and fourth amino acids differ from those of CSF-1.

8. A recombinant baculovirus transfer vector according to claim 3 wherein a NcoI restriction endonuclease recognition site has been introduced at the ATG translational initiation codon of the polyhedrin gene.

9. A recombinant baculovirus transfer vector according to claim 4 wherein the DNA sequence encoding the native polypeptide leader sequence of G-CSF has been deleted from the DNA sequence encoding G-CSF from pP12.

10. A recombinant baculovirus expression vector for the production of biologically active colony-stimulating factors (CSFs) said expression vector comprising a recombinant baculovirus genome and a DNA sequence encoding a CSF under the transcriptional control of a baculovirus promoter.

11. A method for producing a polypeptide with colony stimulating factor (CSF) activity comprising infecting a susceptible host insect cell with a recombinant baculovirus expression vector wherein said expression vector is a recombinant baculovirus genome comprising at least one DNA sequence encoding a CSF under the transcriptional control of a baculovirus promoter, growing said infected insect cells under suitable conditions and recovering said polypeptide from the culture medium.

12. The method of claim 11 wherein said host insect cells are Spodoptera frugiperda cells.

13. The method of claim 11 wherein said baculovirus genome is derived from Autographa californica, Trichoplusia ni, Rachiplusia ou or Galleria mellonella.

14. The method of claim 11 wherein the recombinant baculovirus transfer vector containing a gene encoding a CSF is selected from the group consisting of pAcM1, pAcM2, pAcM3, pAcM4, pAcM5, pAcM6, pAcM9, pAcM10, pAcM11, pAcM12, pAcM13, pAcG1 or pAcG2.

AGTGAGGCTC GGCCCGGGGA AAGTGAAAGT TTGCTTGGGT CCTCTCGGGG CCAGAGCCCG TCCTCCGATC CCAGGACAGC GGTGGGGGCC TCGCCGGGG CGCCCACTCC GCAGCAGCCA
 20 40 60 80 100 120
 GCGAGCGAGC GAGCGAGCGA GGGCGGGCGA CGGCGCCGCG CTGCCCGT ATG ACC GCG CCG GGC GCC GGC GGC TGC CCT CCC ACG ACA TGG CTG GGC
 140 160 180 200 220
 Met Thr Ala Pro Gly Ala Ala Gly Arg Cys Pro Thr Thr Trp Leu Gly
 -32
 TCC CTG CTG TTG TTC TGT CTC TGT GTC AGG AGT ATC ACC GAG GAG GTG TCG GAG TAC TGT AGC CAC ATG ATT GGG AGT GGA CAC CTG CAG TCT
 240 260 280 300 320
 Ser Leu Leu Leu Val Cys Leu Leu Ala Ser Arg Ser Ile Thr Glu Glu Val Ser Glu Tyr Cys Ser His Met Ile Gly Ser Gly His Leu Gln Ser
 1
 CTG CAG CGG CTG ATT GAC AGT CAG ATG GAG ACC TCG TGC CAA ATT ACA TTT GAG TTT GTA GAC CAG GAA CAG TTG AAA GAT CCA GTG TGC TAC CTT AAG
 340 360 380 400 420
 Leu Gln Arg Leu Ile Asp Ser Gln Met Glu Thr Ser Cys Gln Ile Thr Phe Glu Phe Val Asp Gln Glu Leu Lys Asp Pro Val Cys Tyr Leu Lys
 20 INTRON SEQUENCE 40
 AAG GCA TTT CTC CTG GTC CAA TAC ATA ATG GAG GAC ACC ATG CGC TTC AGA GAT AAC ACC CCC AAT GCC ATC GCC ATT GTG CAG CTG CAG GAA CTC TCT
 440 460 480 500 520
 Lys Ala Phe Leu Leu Val Gln Tyr Ile Met Glu Asp Thr Met Arg Phe Arg Asp Asn Thr Pro Asn Ala Ile Ala Ile Val Gln Leu Glu Leu Ser
 * 60 80
 TTG AGG CTG AAG AGC TGC TTC ACC AAG GAT TAT GAA GAG CAT GAC AAG GCC TGC GTC CGA ACT TTC TAT GAG ACA CCT CTC CAG TTG CTG GAG AAG GTC
 540 560 580 600 620
 Leu Arg Leu Lys Ser Cys Phe Thr Lys Asp Tyr Glu Glu His Asp Lys Ala Cys Val Arg Thr Phe Tyr Glu Thr Pro Leu Gln Leu Leu Lys Val
 100
 AAG AAT GTC TTT AAT GAA ACA AAG AAT CTC CTT GAC AAG GAC TGG AAT ATT TTC AGC AAG AAC TGC AAC AAC AGC TTT GCT GAA TGC TCC AGC CAA GGC
 640 660 680 700 720
 Lys Asn Val Phe Asn Glu Thr Lys Asn Leu Leu Asp Lys Asp Trp Asn Ile Phe Ser Lys Asn Cys Asn Asn Ser Phe Ala Glu Cys Ser Ser Gln Gly
 120 140 160

1 / 10

FIG. 1-1

CAT GAG AGG CAG TCC GAG GGA TCC AGC CCG CAG CTC CAG GAG TCT GTC TTC CAC CTG CTG GTG CCC AGT GTC ATC CTG GTC TTG CTG GCC GTC GGA
 His Glu Arg Gln Ser Glu Gly Ser Ser Ser Pro Gln Leu Gln Ser Val Phe His Leu Leu Val Ile Leu Val Leu Leu Ala Val Gly
 160 180 200 220 240 260 280 300 320 340 360 380 400 420 440 460 480 500 520 540 560 580 600 620 640 660 680 700 720 740 760 780 800 820 840 860 880 900 920 940 960 980 1000 1020 1040 1060 1080 1100 1120 1140 1160 1180 1200 1220 1240 1260 1280 1300 1320 1340 1360 1380 1400 1420 1440 1460 1480 1500 1520 1540 1560 1580 1600 1620 1640 1660 1680 1700 1720 1740 1760 1780 1800 1820 1840 1860 1880 1900 1920 1940 1960 1980 2000 2020 2040 2060 2080 2100 2120 2140 2160 2180 2200 2220 2240 2260 2280 2300 2320 2340 2360 2380 2400 2420 2440 2460 2480 2500 2520 2540 2560 2580 2600 2620 2640 2660 2680 2700 2720 2740 2760 2780 2800 2820 2840 2860 2880 2900 2920 2940 2960 2980 3000 3020 3040 3060 3080 3100 3120 3140 3160 3180 3200 3220 3240 3260 3280 3300 3320 3340 3360 3380 3400 3420 3440 3460 3480 3500 3520 3540 3560 3580 3600 3620 3640 3660 3680 3700 3720 3740 3760 3780 3800 3820 3840 3860 3880 3900 3920 3940 3960 3980 4000 4020 4040 4060 4080 4100 4120 4140 4160 4180 4200 4220 4240 4260 4280 4300 4320 4340 4360 4380 4400 4420 4440 4460 4480 4500 4520 4540 4560 4580 4600 4620 4640 4660 4680 4700 4720 4740 4760 4780 4800 4820 4840 4860 4880 4900 4920 4940 4960 4980 5000 5020 5040 5060 5080 5100 5120 5140 5160 5180 5200 5220 5240 5260 5280 5300 5320 5340 5360 5380 5400 5420 5440 5460 5480 5500 5520 5540 5560 5580 5600 5620 5640 5660 5680 5700 5720 5740 5760 5780 5800 5820 5840 5860 5880 5900 5920 5940 5960 5980 6000 6020 6040 6060 6080 6100 6120 6140 6160 6180 6200 6220 6240 6260 6280 6300 6320 6340 6360 6380 6400 6420 6440 6460 6480 6500 6520 6540 6560 6580 6600 6620 6640 6660 6680 6700 6720 6740 6760 6780 6800 6820 6840 6860 6880 6900 6920 6940 6960 6980 7000 7020 7040 7060 7080 7100 7120 7140 7160 7180 7200 7220 7240 7260 7280 7300 7320 7340 7360 7380 7400 7420 7440 7460 7480 7500 7520 7540 7560 7580 7600 7620 7640 7660 7680 7700 7720 7740 7760 7780 7800 7820 7840 7860 7880 7900 7920 7940 7960 7980 8000 8020 8040 8060 8080 8100 8120 8140 8160 8180 8200 8220 8240 8260 8280 8300 8320 8340 8360 8380 8400 8420 8440 8460 8480 8500 8520 8540 8560 8580 8600 8620 8640 8660 8680 8700 8720 8740 8760 8780 8800 8820 8840 8860 8880 8900 8920 8940 8960 8980 9000 9020 9040 9060 9080 9100 9120 9140 9160 9180 9200 9220 9240 9260 9280 9300 9320 9340 9360 9380 9400 9420 9440 9460 9480 9500 9520 9540 9560 9580 9600 9620 9640 9660 9680 9700 9720 9740 9760 9780 9800 9820 9840 9860 9880 9900 9920 9940 9960 9980 10000 10020 10040 10060 10080 10100 10120 10140 10160 10180 10200 10220 10240 10260 10280 10300 10320 10340 10360 10380 10400 10420 10440 10460 10480 10500 10520 10540 10560 10580 10600 10620 10640 10660 10680 10700 10720 10740 10760 10780 10800 10820 10840 10860 10880 10900 10920 10940 10960 10980 11000 11020 11040 11060 11080 11100 11120 11140 11160 11180 11200 11220 11240 11260 11280 11300 11320 11340 11360 11380 11400 11420 11440 11460 11480 11500 11520 11540 11560 11580 11600 11620 11640 11660 11680 11700 11720 11740 11760 11780 11800 11820 11840 11860 11880 11900 11920 11940 11960 11980 12000 12020 12040 12060 12080 12100 12120 12140 12160 12180 12200 12220 12240 12260 12280 12300 12320 12340 12360 12380 12400 12420 12440 12460 12480 12500 12520 12540 12560 12580 12600 12620 12640 12660 12680 12700 12720 12740 12760 12780 12800 12820 12840 12860 12880 12900 12920 12940 12960 12980 13000 13020 13040 13060 13080 13100 13120 13140 13160 13180 13200 13220 13240 13260 13280 13300 13320 13340 13360 13380 13400 13420 13440 13460 13480 13500 13520 13540 13560 13580 13600 13620 13640 13660 13680 13700 13720 13740 13760 13780 13800 13820 13840 13860 13880 13900 13920 13940 13960 13980 14000 14020 14040 14060 14080 14100 14120 14140 14160 14180 14200 14220 14240 14260 14280 14300 14320 14340 14360 14380 14400 14420 14440 14460 14480 14500 14520 14540 14560 14580 14600 14620 14640 14660 14680 14700 14720 14740 14760 14780 14800 14820 14840 14860 14880 14900 14920 14940 14960 14980 15000 15020 15040 15060 15080 15100 15120 15140 15160 15180 15200 15220 15240 15260 15280 15300 15320 15340 15360 15380 15400 15420 15440 15460 15480 15500 15520 15540 15560 15580 15600 15620 15640 15660 15680 15700 15720 15740 15760 15780 15800 15820 15840 15860 15880 15900 15920 15940 15960 15980 16000 16020 16040 16060 16080 16100 16120 16140 16160 16180 16200 16220 16240 16260 16280 16300 16320 16340 16360 16380 16400 16420 16440 16460 16480 16500 16520 16540 16560 16580 16600 16620 16640 16660 16680 16700 16720 16740 16760 16780 16800 16820 16840 16860 16880 16900 16920 16940 16960 16980 17000 17020 17040 17060 17080 17100 17120 17140 17160 17180 17200 17220 17240 17260 17280 17300 17320 17340 17360 17380 17400 17420 17440 17460 17480 17500 17520 17540 17560 17580 17600 17620 17640 17660 17680 17700 17720 17740 17760 17780 17800 17820 17840 17860 17880 17900 17920 17940 17960 17980 18000 18020 18040 18060 18080 18100 18120 18140 18160 18180 18200 18220 18240 18260 18280 18300 18320 18340 18360 18380 18400 18420 18440 18460 18480 18500 18520 18540 18560 18580 18600 18620 18640 18660 18680 18700 18720 18740 18760 18780 18800 18820 18840 18860 18880 18900 18920 18940 18960 18980 19000 19020 19040 19060 19080 19100 19120 19140 19160 19180 19200 19220 19240 19260 19280 19300 19320 19340 19360 19380 19400 19420 19440 19460 19480 19500 19520 19540 19560 19580 19600 19620 19640 19660 19680 19700 19720 19740 19760 19780 19800 19820 19840 19860 19880 19900 19920 19940 19960 19980 20000 20020 20040 20060 20080 20100 20120 20140 20160 20180 20200 20220 20240 20260 20280 20300 20320 20340 20360 20380 20400 20420 20440 20460 20480 20500 20520 20540 20560 20580 20600 20620 20640 20660 20680 20700 20720 20740 20760 20780 20800 20820 20840 20860 20880 20900 20920 20940 20960 20980 21000 21020 21040 21060 21080 21100 21120 21140 21160 21180 21200 21220 21240 21260 21280 21300 21320 21340 21360 21380 21400 21420 21440 21460 21480 21500 21520 21540 21560 21580 21600 21620 21640 21660 21680 21700 21720 21740 21760 21780 21800 21820 21840 21860 21880 21900 21920 21940 21960 21980 22000 22020 22040 22060 22080 22100 22120 22140 22160 22180 22200 22220 22240 22260 22280 22300 22320 22340 22360 22380 22400 22420 22440 22460 22480 22500 22520 22540 22560 22580 22600 22620 22640 22660 22680 22700 22720 22740 22760 22780 22800 22820 22840 22860 22880 22900 22920 22940 22960 22980 23000 23020 23040 23060 23080 23100 23120 23140 23160 23180 23200 23220 23240 23260 23280 23300 23320 23340 23360 23380 23400 23420 23440 23460 23480 23500 23520 23540 23560 23580 23600 23620 23640 23660 23680 23700 23720 23740 23760 23780 23800 23820 23840 23860 23880 23900 23920 23940 23960 23980 24000 24020 24040 24060 24080 24100 24120 24140 24160 24180 24200 24220 24240 24260 24280 24300 24320 24340 24360 24380 24400 24420 24440 24460 24480 24500 24520 24540 24560 24580 24600 24620 24640 24660 24680 24700 24720 24740 24760 24780 24800 24820 24840 24860 24880 24900 24920 24940 24960 24980 25000 25020 25040 25060 25080 25100 25120 25140 25160 25180 25200 25220 25240 25260 25280 25300 25320 25340 25360 25380 25400 25420 25440 25460 25480 25500 25520 25540 25560 25580 25600 25620 25640 25660 25680 25700 25720 25740 25760 25780 25800 25820 25840 25860 25880 25900 25920 25940 25960 25980 26000 26020 26040 26060 26080 26100 26120 26140 26160 26180 26200 26220 26240 26260 26280 26300 26320 26340 26360 26380 26400 26420 26440 26460 26480 26500 26520 26540 26560 26580 26600 26620 26640 26660 26680 26700 26720 26740 26760 26780 26800 26820 26840 26860 26880 26900 26920 26940 26960 26980 27000 27020 27040 27060 27080 27100 27120 27140 27160 27180 27200 27220 27240 27260 27280 27300 27320 27340 27360 27380 27400 27420 27440 27460 27480 27500 27520 27540 27560 27580 27600 27620 27640 27660 27680 27700 27720 27740 27760 27780 27800 27820 27840 27860 27880 27900 27920 27940 27960 27980 28000 28020 28040 28060 28080 28100 28120 28140 28160 28180 28200 28220 28240 28260 28280 28300 28320 28340 28360 28380 28400 28420 28440 28460 28480 28500 28520 28540 28560 28580 28600 28620 28640 28660 28680 28700 28720 28740 28760 28780 28800 28820 28840 28860 28880 28900 28920 28940 28960 28980 29000 29020 29040 29060 29080 29100 29120 29140 29160 29180 29200 29220 29240 29260 29280 29300 29320 29340 29360 29380 29400 29420 29440 29460 29480 29500 29520 29540 29560 29580 29600 29620 29640 29660 29680 29700 29720 29740 29760 29780 29800 29820 29840 29860 29880 29900 29920 29940 29960 29980 30000 30020 30040 30060 30080 30100 30120 30140 30160 30180 30200 30220 30240 30260 30280 30300 30320 30340 30360 30380 30400 30420 30440 30460 30480 30500 30520 30540 30560 30580 30600 30620 30640 30660 30680 30700 30720 30740 30760 30780 30800 30820 30840 30860 30880 30900 30920 30940 30960 30980 31000 31020 31040 31060 31080 31100 31120 31140 31160 31180 31200 31220 31240 31260 31280 31300 31320 31340 31360 31380 31400 31420 31440 31460 31480 31500 31520 31540 31560 31580 31600 31620 31640 31660 31680 31700 31720 31740 31760 31780 31800 31820 31840 31860 31880 31900 31920 31940 31960 31980 32000 32020 32040 32060 32080 32100 32120 32140 32160 32180 32200 32220 32240 32260 32280 32300 32320 32340 32360 32380 32400 32420 32440 32460 32480 32500 32520 32540 32560 32580 32600 32620 32640 32660 32680 32700 32720 32740 32760 32780 32800 32820 32840 32860 32880 32900 32920 32940 32960 32980 33000 33020 33040 33060 33080 33100 33120 33140 33160 33180 33200 33220 33240 33260 33280 33300 33320 33340 33360 33380 33400 33420 33440 33460 33480 33500 33520 33540 33560 33580 33600 33620 33640 33660 33680 33700 33720 33740 33760 33780 33800 33820 33840 33860 33880 33900 33920 33940 33960 33980 34000 34020 34040 34060 34080 34100 34120 34140 34160 34180 34200 34220 34240 34260 34280 34300 34320 34340 34360 34380 34400 34420 34440 34460 34480 34500 34520 34540 34560 34580 34600 34620 34640 34660 34680 34700 34720 34740 34760 34780 34800 34820 34840 34860 34880 34900 34920 34940 34960 34980 35000 35020 35040 35060 35080 35100 35120 35140 35160 35180 35200 35220 35240 35260 35280 35300 35320 35340 35360 35380 35400 35420 35440 35460 35480 35500 35520 35540 35560 35580 35600 35620 35640 35660 35680 35700 35720 35740 35760 35780 35800 35820 35840 35860 35880 35900 35920 35940 35960 35980 36000 36020 36040 36060 36080 36100 36120 36140 36160 36180 36200 36220 36240 36260 36280 36300 36320 36340 36360 36380 36400 36420 36440 36460 36480 36500 36520 36540 36560 36580 36600 36620 36640 36660 36680 36700 36720 36740 36760 36780 36800 36820 36840 36860 36880 36900 36920 36940 36960 36980 37000 37020 37040 37060 37080 37100 37120 37140 37160 37180 37200 37220 37240 37260 37280 37300 37320 37340 37360 37380 37400 37420 37440 37460 37480 37500 37520 37540 37560 37580 37600 37620 37640 37660 37680 37700 37720 37740 37760 37780 37800 37820 37840 37860 37880 37900 37920 37940 37960 37980 38000 38020 38040 38060 38080 38100 38120 38140 38160 38180 38200 38220 38240 38260 38280 38300 38320 38340 38360 38380 38400 38420 38440 38460 38480 38500 38520 38540 38560 38580 38600 38620 38640 38660 38680 38700 38720 38740 38760 38780 38800 38820 38840 38860 38880 38900 38920 38940 38960 38980 39000 39020 39040 39060 39080 39100 39120 39140 39160 39180 39200 39220 39240 39260 39280 39300 39320 39340 39360 39380 39400 39420 39440 39460 39480 39500 39520 39540 39560 39580 39600 39620 39640 39660 39680 39700 39720 39740 39760 39780 39800 39820 39840 39860 39880 39900 39920 39940 39960 39980 40000 40020 40040 40060 40080 40100 40120 40140 40160 40180 40200 40220 40240 40260 40280 40300 40320 40340 40360 40380 40400 40420 40440 40460 40480 40500 40520 40540 40560 40580 40600 40620 40640 40660 40680 40700 40720 40740 40760 40780 40800 40820 40840 40860 40880 40900 40920 40940 40960 40980 41000 41020 41040 41060 41080 41100 41120 41140 41160 41180 41200 41220 41240 41260 41280 41300 41320 41340 41360 41380 41400 41420 41440 41460 41480 41500 41520 41540 41560 41580 41600 41620 41640 41660 41680 41700 41720 41740 41760 41780 41800 41820 41840 41860 41880 41900 41920 41940 41960 41980 42000 42020 42040 42060 42080 42100 42120 42140 42160 42180 42200 42220 42240 42260 42280 42300 42320 42340 42360 42380 42400 42420 42440 42460 4

3 / 10

1 CCCTGGCTGTTGGTCTCTCCCTGGCGAGACGAGAGTATACACGAGGAGGTGTCGGAGTACTGTAGCCACATGATTGGAGTGGACACCTGCAGTCTCTGCAGCGGCTGACAGTC
 LeuLeuLeuValCysLeuLeuAlaSerArgSerIleThrGluGluValSerGluTyrCysSerHisMETIleGlySerGlyHisLeuGlnSerLeuGlnArgLeuIleAspSerGln
 -10 1 20
 121 AGATGGAGACCTCGTGGCCAAATTACATTTGAGTTGTAGACGAGGAACAGTGAAGATCCAGTGTGTACCTTAAGAGGCATTTCCTGGTACAGACATAATGGAGGACACCATGC
 METGluThrSerCysGlnIleThrPheGluPheValAspGlnGluGlnLeuLysAspProValCystyrLeuLysLysAlaPheLeuLeuValGlnAspIleMETGluAspThrMETArg
 40 60 IV
 241 GCTTCAGAGATAACACCCCCCAATGCCATCGCCATTGTGCAGTGCAGGAACCTCTCTTTGAGGCTGAAGAGCTGCTTCAACAAGGATTATGAAGAGCATGACAAGCGCTCGTCCGAACCTT
 PheArgAspAsnThrProAsnAlaIleAlaIleValGlnLeuSerLeuArgLeuLysSerCysPheThrLysAspTyrGluGluHisAspLysAlaCysValArgThrPhe
 80 100
 361 TCTATGAGACACCTCTCCAGTTGCTGGAGAGAGTCAAGAATGTCTTTAATGAACAACAAGAAATCTCCTTGACAAGGACTGGAAATATTTTCAGCAAGAACTGCAACAACAGCTTTGCTGAAT
 TyrGluThrProLeuGlnLeuLeuGluLysValLysAsnValPheAsnGluThrLysAsnLeuLeuAspLysAspTrpAsnIlePheSerLysAsnCysAsnAsnSerPheAlaGluCys
 Y 120 140
 481 GCTCCAGGCAAGATGTGGTGACCAAGCCTGATTGCCAAGTCCCTGTACCCCAAGCCATCCCTAGCAGTGACCGGCGCTGTCTCCCTCATCAGCGCCCTCGCCCTCCATGGCCCTG
 SerSerGlnAspValValThrLysProAspCysAsnCysLeuTyrProLysAlaIleProSerSerAspProAlaSerValSerProHisGlnProLeuAlaProSerMETAlaProVal
 160 180
 601 TGGCTGGCTTGACCTGGGAGGACTCTGAGGGAAGTCAAGGCGAGTCCCTCTGCTGGTGTAGCAGCCCTGCACACAGTGGATCCAGSCAGTGCACAGCGGCGCACCCAGGAGCACCT
 AlaGlyLeuThrTrpGluAspSerGluGlyThrGluGlySerSerLeuLeuProGlyGluGlnProLeuHisThrValAspProGlySerAlaLysGlnArgProProArgSerThrCys
 200 220
 721 GCCAGAGCTTTGAGCGCCAGAGACCCAGTTGTCAAGGACACACCATCGGTGGCTCACACAGCCTCGCCCTCTGTGGGGCGCTTCAACCCCGGATGGAGGATATCTTGACTCTG
 GlnSerPheGluProProGluThrProValValLysAspSerThrIleGlyGlySerProGlnProArgProSerValGlyAlaPheAsnProGlyMETGluAspIleLeuAspSerAla
 240 260
 841 CAATGGGCACTAATTGGGTCCAGAGAGAGCCCTGTGGAGAGCCAGTGAGATTCCCGTACCCCAAGGACAGAGCTTTCCCTCCAGGCCAGGAGGGGCGAGCATGCAGACAGAGCCCG
 METGlyThrAsnTrpValProGluGluAlaSerGlyGluAlaSerGluIleProValProGlnGlyThrGluLeuSerProSerArgProGlyGlyGlySerMETGlnThrGluProAla
 280 300

FIG. 2-1

SUBSTITUTE SHEET

961 CCAGACCCAGCAACTTCTCCTCAGCATCTTCTCCACTCCCTGCATCAGCAAAGGGCCAAGCGGACAGATGTAACTGGTACAGCCTTGCCCAGGGTTGGCCCCCGTGTAGGCCACTGGCC
ArgProSerAsnPheLeuSerAlaSerProLeuProAlaSerProLeuProAlaSerValThrGlyThrAlaLeuProArgValGlyProValArgProThrGlyGln
320 340

1081 AGACTGGGAATCACACCCCCAGAAGACAGACCATTCCATCTGCCCTGCCTCAGAGACCCCGGAGCCAGGCTCTCCAGGATCTCATCTACCTGCCCCCCCAGGGCCTCAGCAACCCCTCCA
AspTrpAsnHisThrProGlnLysThrAspHisProSerAlaLeuLeuA-GaspProProGluProGlySerProArgIleSerSerLeuArgProGlnGlyLeuSerAsnProSerThr
360 380

1201 CCTCTCTGCTCAGCCACAGCTTTCAGAACCCACTCTCCTCGGACGCGTGTGCCCTTGGGAGCTGGAGGGCAGGAGGAGCACAGGGATCGAGAGGAGGCCCGCAGAGCCAGGACGAGAAGGAG
LeuSerAlaGlnProGlnLeuSerArgSerHisSerSerGlySerValLeuProLeuGlyGluLeuGluGlyArgSerThrArgAspArgSerProAlaGluProGluGlyGly
400 420

1321 GACCAAGAGTAGGGGGCAGCCAGGCCCTGCCCGCTTTTAACTCGTTCCCTTGTACTGACACAGGCCCATCAGAGSCAGTCCGAGGGATCTCTCAGCCCGGAGCTCCAGGAGTCTGTCT
ProAlaSerGluGlyAlaAlaArgProLeuProArgPheAsnSerValProLeuThrAspThrGlyHisGluArgGlnSerGluGlySerSerSerProGlnLeuGlnGluSerValPhe
440 460

1441 TCCACCTGCTGGTGCCAGTGTCATCCTGGTCTTGTCTGGCCGTCGGAGGCCCTTGTCTTACAGGTGGAGGGCGGAGGCCATCAAGAGCCCTCAGAGAGCGGATTTCTCCCTGGAGCAAC
HisLeuLeuValProSerValIleLeuValLeuLeuAlaValGlyGlyLeuLeuPheTyArgTrpArgArgArgSerHisGlnGluProGlnArgAlaAspSerProLeuGluGlnPro
VII 480 500

1561 CAGAGGCGAGCCCTGACTCAGGATGACAGACAGGTGGAACTGCCAGTGTAGAGGGAATTCTAAGACCCCTCACCATCCTGGACACACTCGTTTGTCAATGTCCTCTGAAAAATGTGAC
GluGlySerProLeuThrGlnAspAspArgGlnValGluLeuProVal
520

1681 GCCAGGCCCGGACACAGTACTCCAGATGTTGTCTGACCAAGCTCAGAGAGAGTACAGTGGGACTGTTACCTTCCCTTCATATGGACAGATATTCTTCTATTGTCGAGATTAAGATTGCA
TAGTTTTTTTCTTAAACAACCTGCATCATCTGTGTGTCATATGTTGAGCCTGTGGTCTATTAAAACCCCTAGTTCCATTTCCTATAAACTTCTGTCAGCCAGACCATCTCTACCTGTAC
1921 TTGGACAACCTTAACCTTTTAAACAAGTCAGATTATGTTTACCTTTGTTAAAGCCACCTTGTTGCTGCCATCCTGAACCTACTAGATTGTGTAATCTCTAATCTGTCA
TCTCCGTAGCCCCCTCCGATGTGGCTTCTCGACATTGATGAGTGCCTGTGTTCTTGGCCCATGTGTGATGTAGCTGTGACCTATTGTTCTCTCACCCCTGCCCCCGCCCAACCC
2161 CAGCTGGCCCAACCTCTTCCCTTCCCAACCCCAAGCCCAAGCCCATCAGGAGCCCTTCCCTGGCTTCTCCACAACCTTCTGACTGCTCTTTTTCAGTGTGCCCCCTCTCTCTCTTTT
TATTGGCTATAGTATATCAATTGC

FIG. 2-2

5 / 10

FIG. 3

1 AAAAAACAGCCCGGAGCCTGCAGCCAGCCCCACCCAGACCCATGGCTGGACCTGCCACC
MetAlaGlyProAlaThr
-30

61 CAGAGCCCCATGAAGCTGATGGCCCTGCAGCTGCTGCTGTGGCACAGTGCACCTCTGGACA
GlnSerProMetLysLeuMetAlaLeuGlnLeuLeuLeuTrpHisSerAlaLeuTrpThr
-20 -10

121 GTGCAGGAAGCCACCCCTGGGCCCTGCCAGCTCCCTGCCCCAGAGCTTCCTGCTCAAG
ValGlnGluAlaThrProLeuGlyProAlaSerSerLeuProGlnSerPheLeuLeuLys
-1 1 10

181 TGCTTAGAGCAAGTGAGGAAGATCCAGGGCGATGGCGCAGCGCTCCAGGAGAAGCTGTGT
CysLeuGluGlnValArgLysIleGlnGlyAspGlyAlaAlaLeuGlnGluLysLeuCys
20 30

241 GCCACCTACAAGCTGTGCCACCCGAGGAGCTGGTGCTGCTCGGACACTCTCTGGGCATC
AlaThrTyrLysLeuCysHisProGluGluLeuValLeuLeuGlyHisSerLeuGlyIle
40 50

301 CCCTGGGCTCCCCTGAGCAGCTGCCCCAGCCAGGCCCTGCAGCTGGCAGGCTGCTTGAGC
ProTrpAlaProLeuSerSerCysProSerGlnAlaLeuGlnLeuAlaGlyCysLeuSer
60 70

361 CAACTCCATAGCGGCCTTTTCCTCTACCAGGGGCTCCTGCAGGCCCTGGAAGGGATCTCC
GlnLeuHisSerGlyLeuPheLeuTyrGlnGlyLeuLeuGlnAlaLeuGluGlyIleSer
80 90

421 CCCGAGTTGGGTCCCACCTTGGACACACTGCAGCTGGACGTCGCCGACTTTGCCACCACC
ProGluLeuGlyProThrLeuAspThrLeuGlnLeuAspValAlaAspPheAlaThrThr
100 110

481 ATCTGGCAGCAGATGGAAGAACTGGGAATGGCCCCCTGCCCTGCAGCCCACCCAGGGTGCC
IleTrpGlnGlnMetGluGluLeuGlyMetAlaProAlaLeuGlnProThrGlnGlyAla
120 130

541 ATGCCGGCCTTCGCCTCTGCTTTCCAGCGCCGGGCAGGAGGGGTCTAGTTGCCTCCCAT
MetProAlaPheAlaSerAlaPheGlnArgArgAlaGlyGlyValLeuValAlaSerHis
140 150

601 CTGCAGAGCTTCCTGGAGGTGTCGTACCGGTTCTACGCCACCTTGCCAGCCCTGAGCC
LeuGlnSerPheLeuGluValSerTyrArgValLeuArgHisLeuAlaGlnPro
160 170

661 AAGCCCTCCCCATCCCATGTATTTATCTCTATTTAATATTTATGTCTATTTAAGCCTCAT
721 ATTTAAAGACAGGGAAGAGCAGAACGGAGCCCCAGGCCTCTGTGTCCTTCCCTGCATTTCT
781 TGAGTTTCATTCTCCTGCCTGTAGCAGTGAGAAAAAGCTCCTGTCTCTCCCATCCCTGGA
841 CTGGGAGGTAGATAGGTAAATACCAAGTATTTATTACTATGACTGCTCCCCAGCCCTGGC
901 TCTGCAATGGGCACTGGGATGAGCCGCTGTGAGCCCCCTGGTCCTGAGGGTCCCCACCTGG
961 GACCCTTGAGAGTATCAGGTCTCCACGTTGGGAGACAAGAAATCCCTGTTTAATATTTAA
1021 ACAGCAGTGTTCCCCATCTGGGTCTTGACCCCTCACTCTGGCCTCAGCCGACTGCACA
1081 GCGGCCCTGCATCCCTTGGCTGTGAGGCCCTGGACAAGCAGAGGTGGCCAGAGCTGG
1141 GAGGCATGGCCCTGGGGTCCCACGAATTTGCTGGGGAATCTCGTTTTTCTTCTTAAGACT
1201 TTTGGGACATGGTTTGAATCCCGAACATCACCAGCGTGTCTCTGTTTTTCTGGGTGGCC
1261 TCGGGACACCTGCCCTGCCCCACGAGGGTCAGGACTGTGACTCTTTTATAGGGCCAGGCA
1321 GGTGCTTGGACATTTGCCTTGGCTGGACGGGGACTGGGGATGTGGGAGGGAGCAGACAGGA
1381 GGAATCATGTACGGCCTGTGTGTGAAAGGAAGCTCCACTGTACCCCTCCACCTCTTCACC
1441 CCCCACCTCACCAGTGTCCCCTCCACTGTCACATTGTAAGTGAAGTTAGGATAATAAGAT
1501 GTTTGCCTCCAAAAAAAAAAAAAAAAA

SUBSTITUTE SHEET

6 / 10





FIG. 4

BACULOVIRUS TRANSFER VECTORS

SUBSTITUTE SHEET

FIG. 6



9 / 10

FIG. 7

PLASMID	CODING SEQUENCE	CONSTRUCTION OF PLASMID	EXPRESSION (RIA ASSAY) SMALL SCALE INFECTIONS
pAcM1	VAL 224	IN pAc611	7500 U/mL
pAcM2	SER 158	IN pAc611	75,000
pAcM3	SER 158	REMOVE POLYLINKER AND CSF-1 LEADER	320,000
pAcM4	GLY 150	SMA I SITE IN CSF-1, PUT IN pAcC1	500,000 (1,200,000 SCALED-UP)
pAcM5	VAL 522	IN pAc610	24,000
pAcM6	VAL 522	IN pAcM4	100,000
pAcM9	GLY 150, GLN 122, GLN 140	IN pAcM4	24,000
pAcM10	GLY 150	IN pAcC3	500,000
pAcM11	VAL 522, GLN 122, GLN 140	IN pAcM4	13,000
pAcM12	PRO 221	IN pAcM4	>80,000
pAcM13	PRO 221, GLN 122, GLN 140	IN pAcM4	>80,000

SUBSTITUTE SHEET

10 / 10




FIG. 8

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 88/02445

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) * According to International Patent Classification (IPC) or to both National Classification and IPC IPC ⁴ : C 12 N 15/00																				
II. FIELDS SEARCHED <div style="text-align: right; font-size: small;">Minimum Documentation Searched ⁷</div> <table style="width: 100%; border: none;"> <tr> <td style="width: 25%; border: none;">Classification System</td> <td style="border: none;">Classification Symbols</td> </tr> <tr> <td style="border: none; padding-top: 10px;">IPC⁴</td> <td style="border: none; padding-top: 10px;">C 12 N</td> </tr> </table> <div style="text-align: center; font-size: x-small; margin-top: 10px;"> Documentation Searched other than Minimum Documentation to the extent that such Documents are Included in the Fields Searched ⁸ </div>			Classification System	Classification Symbols	IPC ⁴	C 12 N														
Classification System	Classification Symbols																			
IPC ⁴	C 12 N																			
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹ <table border="1" style="width: 100%; border-collapse: collapse; font-size: small;"> <thead> <tr> <th style="width: 10%;">Category ¹⁰</th> <th style="width: 60%;">Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²</th> <th style="width: 30%;">Relevant to Claim No. ¹³</th> </tr> </thead> <tbody> <tr> <td style="text-align: center; vertical-align: top;">Y</td> <td>WO, A, 86/00639 (SANDOZ AG) 30 January 1986 see claims --</td> <td style="text-align: center; vertical-align: top;">1, 2, 10, 11</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">Y</td> <td>EP, A, 0127839 (THE TEXAS A&M UNIVERSITY SYSTEM) 12 December 1984 see claims cited in the application --</td> <td style="text-align: center; vertical-align: top;">1, 2, 10, 11</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">Y</td> <td>Proc. Natl. Acad. Sci, USA, volume 82, no. 24, December 1985, (Washington DC, US), G.E. Smith et al.: "Modification and secretion of human interleukin 2 produced in insect cells by a baculo-virus expression vector", pages 8404-8408 see the whole article --</td> <td style="text-align: center; vertical-align: top;">1, 2, 10, 11</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">P, X</td> <td>EP, A, 0272779 (CETUS CORP.) 29 June 1988 see page 10, lines 3-6; claims --</td> <td style="text-align: center; vertical-align: top;">10, 11</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">P, X</td> <td>Gene, volume 58, nos. 2, 3, 1987, Elsevier Science Publishers B.V. (Biomedical ./.)</td> <td style="text-align: center; vertical-align: top;">1, 2, 10, 11</td> </tr> </tbody> </table>			Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³	Y	WO, A, 86/00639 (SANDOZ AG) 30 January 1986 see claims --	1, 2, 10, 11	Y	EP, A, 0127839 (THE TEXAS A&M UNIVERSITY SYSTEM) 12 December 1984 see claims cited in the application --	1, 2, 10, 11	Y	Proc. Natl. Acad. Sci, USA, volume 82, no. 24, December 1985, (Washington DC, US), G.E. Smith et al.: "Modification and secretion of human interleukin 2 produced in insect cells by a baculo-virus expression vector", pages 8404-8408 see the whole article --	1, 2, 10, 11	P, X	EP, A, 0272779 (CETUS CORP.) 29 June 1988 see page 10, lines 3-6; claims --	10, 11	P, X	Gene, volume 58, nos. 2, 3, 1987, Elsevier Science Publishers B.V. (Biomedical ./.)	1, 2, 10, 11
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³																		
Y	WO, A, 86/00639 (SANDOZ AG) 30 January 1986 see claims --	1, 2, 10, 11																		
Y	EP, A, 0127839 (THE TEXAS A&M UNIVERSITY SYSTEM) 12 December 1984 see claims cited in the application --	1, 2, 10, 11																		
Y	Proc. Natl. Acad. Sci, USA, volume 82, no. 24, December 1985, (Washington DC, US), G.E. Smith et al.: "Modification and secretion of human interleukin 2 produced in insect cells by a baculo-virus expression vector", pages 8404-8408 see the whole article --	1, 2, 10, 11																		
P, X	EP, A, 0272779 (CETUS CORP.) 29 June 1988 see page 10, lines 3-6; claims --	10, 11																		
P, X	Gene, volume 58, nos. 2, 3, 1987, Elsevier Science Publishers B.V. (Biomedical ./.)	1, 2, 10, 11																		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"G" document member of the same patent family</p> </div> </div>																				
IV. CERTIFICATION <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none; vertical-align: top;"> Date of the Actual Completion of the International Search <div style="text-align: center; font-size: large;">27th October 1988</div> International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div> </td> <td style="width: 50%; border: none; vertical-align: top;"> Date of Mailing of this International Search Report <div style="text-align: center; font-size: large;">28 NOV 1988</div> Signature of Authorized Officer <div style="text-align: center;"> P.C.G. VAN DER PUTTEN </div> </td> </tr> </table>			Date of the Actual Completion of the International Search <div style="text-align: center; font-size: large;">27th October 1988</div> International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div>	Date of Mailing of this International Search Report <div style="text-align: center; font-size: large;">28 NOV 1988</div> Signature of Authorized Officer <div style="text-align: center;"> P.C.G. VAN DER PUTTEN </div>																
Date of the Actual Completion of the International Search <div style="text-align: center; font-size: large;">27th October 1988</div> International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div>	Date of Mailing of this International Search Report <div style="text-align: center; font-size: large;">28 NOV 1988</div> Signature of Authorized Officer <div style="text-align: center;"> P.C.G. VAN DER PUTTEN </div>																			

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
	Division), (Amsterdam, NL), A. Miyajima et al.: "Use of the silkworm, Bombyx mori, and an insect baculovirus vector for high-level expression and secretion of bio- logically active mouse interleukin-3", pages 273-281 see the whole article --	
P,X	Biotechnology, volume 6, January 1988, (New York, US), V.A. Luckow et al.: "Trends in the development of baculovirus expression vectors", pages 47-55 see table 1 -----	1,2,10-13

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

US 8802445
SA 23618

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on 15/11/88
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A- 8600639	30-01-86	AU-A- 4545685	10-02-86
		EP-A- 0188479	30-07-86
		JP-T- 61502682	20-11-86
EP-A- 0127839	12-12-84	AU-A- 2871784	29-11-84
		JP-A- 60037988	27-02-85
		CA-A- 1222213	26-05-87
		US-A- 4745051	17-05-88
EP-A- 0272779	29-06-88	AU-A- 8007087	28-04-88
		WO-A- 8803173	05-05-88